

RECONSTRUCTION OF SIGNALS FROM MAGNITUDES OF REDUNDANT REPRESENTATIONS: THE COMPLEX CASE

RADU BALAN

ABSTRACT. This paper is concerned with the question of reconstructing a vector in a finite-dimensional complex Hilbert space when only the magnitudes of the coefficients of the vector under a redundant linear map are known. We present new invertibility results as well an iterative algorithm that finds the least-square solution and is robust in the presence of noise. We analyze its numerical performance by comparing it to the Cramer-Rao lower bound.

1. INTRODUCTION

This paper is concerned with the question of reconstructing a vector x in a finite-dimensional *complex* Hilbert space H when only the magnitudes of the coefficients of the vector under a redundant linear map are known.

Specifically our problem is to reconstruct $x \in H$ up to a global phase factor from the magnitudes $\{|\langle x, f_k \rangle|, 1 \leq k \leq m\}$ where $\{f_1, \dots, f_m\}$ is a frame (complete system) for H . This paper is the companion paper to the arXiv submission [9].

A previous paper [4] described the importance of this problem to signal processing, in particular to the analysis of speech. Of particular interest is the case when the coefficients are obtained from a Windowed Fourier Transform (also known as Short-Time Fourier Transform), or an Undecimated Wavelet Transform (in audio and image signal processing). While [4] presents some necessary and sufficient conditions for reconstruction, the general problem of finding fast/efficient algorithms is still open. In [8] we describe one solution in the case of STFT coefficients. The authors of [57] propose a novel algorithm adapted to compactly supported signals and FFT transforms that uses individual signal spectral powers and of two additional interferences between signals.

For vectors in real Hilbert spaces, the reconstruction problem is easily shown to be equivalent to a combinatorial problem. In [5] this problem is further proved to be equivalent to a (nonconvex) optimization problem.

A different approach (which we called the "algebraic approach") was proposed in [7]. While it applies to both real and complex cases, noiseless and noisy cases, the approach requires solving a linear system of size exponentially in space dimension. The algebraic approach mentioned earlier generalizes the approach in [6] where reconstruction is performed with complexity $O(n^2)$ (plus computation of the principal eigenvector for a matrix of size n) but requiring $m = O(n^2)$ frame vectors.

Recently the authors of [20] developed a convex optimization algorithm (PhaseLift) and proved its ability to perform exact reconstruction in the absence of noise, as well as its

stability under noise conditions. In a separate paper, [21], the authors developed further a similar algorithm in the case of windowed DFT transforms. The original requirement of $m = O(n \log(n))$ vectors has been relaxed to $m = O(n)$ in [22]. Similar convex optimization solutions have been proposed by the authors of [29] and [61]. Additionally, [36] studied the duality gap in this approach and obtained a necessary and sufficient condition for the existence of a dual certificate.

While writing this paper, we have become aware of [11] where the authors obtained similar results to the injectivity criteria presented here, as well as to the Cramer-Rao Lower Bound derived in this paper. We will comment more later in the paper. We also acknowledge the paper [30] where some stability bounds have been obtained in the real case. The real case of the stability bounds obtained here are presented in a separate paper [10] together with additional results for the real case.

The organization of the paper is as follows. In section 2 we define the problem explicitly. In section 3 we describe new analysis results. Specifically we analysis in more details spaces of symmetric operators of constrained signature, and then we shows how they are related to the phaseless recovery problem. Our results are canonical, meaning that are independent to a particular choice of basis. In section 4 we establish two robustness results: bi-Lipschitzianity of the nonlinear analysis map, and the Cramer-Rao lower bound (CRLB). In section 5 we present a new reconstruction algorithm based on the Least-Square method. We also obtain robustness bounds to noise. Its performance is analyzed in section 6 and is compared to the CRLB. Section Section 7 contains conclusions and is followed by references.

2. BACKGROUND

Let H be an n -dimensional complex Hilbert space (such as $\mathbb{C}^n$ or $C^{p_1 \times p_2}$) with scalar product $\langle, \rangle$ linear in the first term and antilinear in the second term and a *conjugation* $c : H \rightarrow H$ (see e.g. [35]; conjugation is an antilinear transformation thatt squares to identity). Let $\mathbb{F} = \{f_1, \dots, f_m\}$ be a spanning set of m vectors in H . In finite dimension (as it is the case here) such a set forms a *frame*. In the infinite dimensional case, the concept of frame involves a stronger property than completeness (see for instance [23]). We review additional terminology and properties which remain still true in the infinite dimensional setting. The set $\mathbb{F}$ is frame if and only if there are two positive constants $0 < A \leq B < \infty$ (called frame bounds) so that

$$A\|x\|^2 \leq \sum_{k=1}^m |\langle x, f_k \rangle|^2 \leq B\|x\|^2$$

When we can choose $A = B$ the frame is said *tight*. For $A = B = 1$ the frame is called *Parseval*. A set of vectors $\mathbb{F}$ of the n -dimensional Hilbert space H is said to have *full spark* if any subset of n vectors is linearly independent (see [2] for a full discussion of such frames).

2.1. Problem Definition and Notations. For a vector $x \in H$, the collection of coefficients $\{\langle x, f_k \rangle, 1 \leq k \leq m\}$ represents the analysis of vector x given by the frame $\mathbb{F}$. In H we

consider the following equivalence relation:

$$(2.1) \quad x, y \in H, \quad x \sim y \text{ iff } y = zx \text{ for some scalar } z \text{ with } |z| = 1$$

Note $z = e^{i\varphi}$ for some real number φ . Let $\hat{H} = H / \sim$ be the set of classes of equivalence induced by this relation. Thus $\hat{x} = \{e^{i\alpha}x, 0 \leq \alpha < 2\pi\}$. The analysis map induces the following nonlinear map

$$(2.2) \quad \alpha : \hat{H} \rightarrow (\mathbb{R}^+)^m, \quad \alpha(\hat{x}) = (|\langle x, f_k \rangle|^2)_{1 \leq k \leq m}$$

where $\mathbb{R}^+ = \{x, x \in \mathbb{R}, x \geq 0\}$ is the set of nonnegative real numbers. In [9] we studied when the nonlinear map α is injective, mostly in the real case, and we provided some necessary conditions of injectivity in the complex case. We review these results below. In this paper we obtain additional injectivity results. We then concentrate on the additive white Gaussian noise model

$$(2.3) \quad y = \alpha(x) + \nu, \quad \nu \sim \mathbb{N}(0, \sigma^2)$$

We describe an algorithm (the Iterative Regularized Least-Square algorithm) to solve the estimation problem. We prove some convergence results and we study its performance in the noisy case. We shall derive the Cramer-Rao Lower Bound (CRLB) for this model and compare the algorithm performance to this bound.

We describe several objects that will be used in this paper.

The set $B(H)$ denotes the set of linear operators on H . In $B(H)$ for any $1 \leq p \leq \infty$, the p -norm of $T \in B(H)$ denoted $\|T\|_p$ is given by the p -norm of its set of singular eigenvalues. In particular for $p = 1$, the 1-norm $\|T\|_1$ is called the nuclear norm of T ; for $p = 2$, the 2-norm $\|T\|_2 = \sqrt{\text{tr}(T^*T)}$ is the Frobenius norm of T ; for $p = \infty$, ∞ -norm $\|T\|_\infty$ is the same as the operator norm of T on H simply denoted $\|T\|$.

For each frame vector f_k we denote by F_k its associated rank-1 operator

$$(2.4) \quad F_k : H \rightarrow H, \quad F_k(x) = \langle x, f_k \rangle f_k$$

In general the *associated rank-1 operator* to a vector $x \in H$ is the operator $X : H \rightarrow H$, $X(v) = \langle v, x \rangle x$. Note this is a slight misnomer: X has *at most* rank one. Specifically, X has rank one if and only if $x \neq 0$; otherwise X has rank zero. One can think of X as simply the $n \times n$ matrix xx^* , where x^* denotes the transpose and conjugate of x . However, as much as possible, we plan to use canonical notations (that is, basis independent).

For any two vectors $u, v \in H$ we define their *symmetric outer product* denoted $\llbracket u, v \rrbracket$ by

$$(2.5) \quad \llbracket u, v \rrbracket : H \rightarrow H, \quad \llbracket u, v \rrbracket(x) = \frac{1}{2}(\langle x, u \rangle v + \langle x, v \rangle u)$$

One can think of $\llbracket u, v \rrbracket$ as the $n \times n$ symmetric matrix $\frac{1}{2}(vu^* + uv^*)$. Note the rank-1 associated to a vector x can be written as $\llbracket x, x \rrbracket$. Thus $F_k = \llbracket f_k, f_k \rrbracket$. Note also $\llbracket u, v \rrbracket$ is $\mathbb{R}$ -bilinear but it is not $\mathbb{C}$ -(bi)linear. Furthermore $\llbracket u, v \rrbracket = \llbracket v, u \rrbracket$.

The nonlinear map α induces a linear map $\mathcal{A}$ on the set $B(H)$ of bounded operators on H :

$$(2.6) \quad \mathcal{A} : B(H) \rightarrow \mathbb{C}^m, \quad (\mathcal{A}(T))_k = \langle T f_k, f_k \rangle = \text{tr}\{T F_k\}$$

In general, for two operators $T, S \in B(H)$ we denote $\langle T, S \rangle_{B(H)} = \text{tr}\{TS^*\}$ their Hilbert-Schmidt scalar product, where S^* is the adjoint of S and tr denotes the trace. Note the scalar product is basis independent, but it depends of the underlying Hilbert space structure. Thus $(\mathcal{A}(T))_k = \langle T, F_k \rangle_{B(H)}$. Also note $\alpha(x) = \mathcal{A}(X)$ where $X = \langle \cdot, x \rangle x$ is the rank-1 associated to x . When no danger of confusion we drop the index $B(H)$ from the scalar product notation.

Let $\text{Sym}(H)$ denote set of self-adjoint operators on H , $\text{Sym}(H) = \{T \in B(H) , T^* = T\}$. We denote by $\mathcal{S}^{p,q}$ or $\mathcal{S}^{p,q}(H)$, the set of self-adjoint operators on H that have at most p positive eigenvalues and at most q negative eigenvalues:

$$(2.7) \quad \begin{aligned} \mathcal{S}^{p,q} = & \{T \in \text{Sym}(H) , Sp(T) = \{\lambda_1, \dots, \lambda_n\} , \\ & \lambda_1 \geq \dots \geq \lambda_q \geq 0 = \lambda_{n-p-q} = \dots = \lambda_{n-q} \geq \lambda_{n-q+1} \geq \dots \geq \lambda_n\} \end{aligned}$$

where $Sp(T)$ denotes the spectrum of T (i.e. the set of its eigenvalues). Notice $\mathcal{S}^{p,q}$ is not a linear space, but instead it is a cone in $B(H)$. This cone property is key in deriving robustness and stability bounds later on.

We denote by $\lambda_{\max}(T)$ the largest eigenvalue of T and by $\lambda_{\min}(T)$ the smallest eigenvalue of T . In particular we are interested in $\mathcal{S}^{1,0}$ and $\mathcal{S}^{1,1}$:

$$(2.8) \quad \mathcal{S}^{1,0} = \{T \in \text{Sym}(H) , \text{rank}(T) \leq 1, \lambda_{\min}(T) = 0\}$$

$$(2.9) \quad \begin{aligned} \mathcal{S}^{1,1} = & \{T \in \text{Sym}(H) , \text{rank}(T) \leq 2, Sp(T) = \{\lambda_{\max}(T), 0^{(n-2)}, \lambda_{\min}(T)\}, \\ & \lambda_{\max}(T) \geq 0 \geq \lambda_{\min}(T)\} \end{aligned}$$

Note the following obvious inclusions

$$(2.10) \quad \{0\} \subset \mathcal{S}^{1,0} \subset \mathcal{S}^{1,1} \subset \text{Sym}(H) , \quad \{0\} \subset \mathcal{S}^{0,1} \subset \mathcal{S}^{1,1} \subset \text{Sym}(H)$$

We denote by $\mathring{\mathcal{S}}^{p,q}$ the subset of $\mathcal{S}^{p,q}$ of selfadjoint operators that have rank $p+q$, hence exactly p strictly positive eigenvalues and q strictly negative eigenvalues. Thus

$$(2.11) \quad \mathcal{S}^{p,q} = \mathring{\mathcal{S}}^{p,q} \cup \mathcal{S}^{p-1,q} \cup \mathcal{S}^{p,q-1}$$

In particular

$$(2.12) \quad \mathcal{S}^{1,1} = \mathring{\mathcal{S}}^{1,1} \cup \mathcal{S}^{1,0} \cup \mathcal{S}^{0,1} , \quad \mathcal{S}^{1,0} = \mathring{\mathcal{S}}^{1,0} \cup \{0\}$$

A more detailed analysis of these sets is presented in subsection 3.1.

Next we describe the realification of the Hilbert space H . To do so canonically we need to fix a conjugation $c : H \rightarrow H$. To the complex Hilbert space H with conjugation c we associate its $2n$ -dimensional real vector space $H_{\mathbb{R}}$ subset of $H \times H$ built from vectors $v_R = \frac{1}{2}(v + c(v))$ and $v_I = \frac{1}{2i}(v - c(v))$ as follows:

$$(2.13) \quad H_{\mathbb{R}} = \left\{ \left(\frac{1}{2}(v + c(v)), \frac{1}{2i}(v - c(v)) \right) , v \in H \right\}$$

Thus $H_{\mathbb{R}}$ is the image of H through the $\mathbb{R}$ -linear map,

$$(2.14) \quad \mathbf{j} : H \rightarrow H \times H , \quad \mathbf{j}(v) = \left(\frac{1}{2}(v + c(v)), \frac{1}{2i}(v - c(v)) \right)$$

Note $\mathbf{j}$ is injective and when restricted to $\mathbf{j} : H \rightarrow H_{\mathbb{R}}$ it becomes an $\mathbb{R}$ - isomorphism. Its inverse is given by

$$(2.15) \quad \mathbf{j}^{-1} : H_{\mathbb{R}} \rightarrow H, \quad \mathbf{j}^{-1}(u, v) = u + iv$$

Let J denote the linear map defined by

$$(2.16) \quad J : H \times H \rightarrow H \times H, \quad J(v, w) = (-w, v)$$

Note it is conjugate to the multiplication by i in H :

$$(2.17) \quad J : H_{\mathbb{R}} \rightarrow H_{\mathbb{R}}, \quad J(\mathbf{j}(v)) = \mathbf{j}(iv)$$

Hence $H_{\mathbb{R}}$ is J invariant. In $H_{\mathbb{R}}$ the induced scalar product is given by

$$(2.18) \quad \langle \mathbf{j}(v), \mathbf{j}(w) \rangle := \left\langle \frac{1}{2}(v + c(v)), \frac{1}{2}(w + c(w)) \right\rangle + \left\langle \frac{1}{2i}(v - ic(v)), \frac{1}{2i}(w - c(w)) \right\rangle = \text{real}(\langle v, w \rangle)$$

We denote by $\langle v, w \rangle_{\mathbb{R}} = \text{real}(\langle v, w \rangle)$ the $\mathbb{R}$ -linear inner product on H . Note

$$(2.19) \quad \begin{aligned} \langle u, v \rangle &= \text{real}(\langle u, v \rangle) + i \text{imag}(\langle u, v \rangle) = \langle u, v \rangle_{\mathbb{R}} - i \langle iu, v \rangle_{\mathbb{R}} \\ &= \langle u, v \rangle_{\mathbb{R}} + i \langle u, iv \rangle_{\mathbb{R}} = \langle \mathbf{j}(u), \mathbf{j}(v) \rangle + i \langle \mathbf{j}(u), J\mathbf{j}(v) \rangle \end{aligned}$$

If $\{e_1, \dots, e_n\}$ is an orthonormal basis in H , then $\{\mathbf{j}(e_1), \dots, \mathbf{j}(e_n), J\mathbf{j}(e_1), \dots, J\mathbf{j}(e_n)\}$ is an orthonormal basis in $H_{\mathbb{R}}$. Which shoes the real dimension of $H_{\mathbb{R}}$ is $2n$, $\dim_{\mathbb{R}} H_{\mathbb{R}} = 2n$.

On $H \times H$ there are two inner product structures:

$$(2.20) \quad \langle (x, y), (u, v) \rangle_{\mathbb{C}} = \langle x, u \rangle + \langle y, v \rangle$$

$$(2.21) \quad \langle (x, y), (u, v) \rangle_{\mathbb{R}} = \frac{1}{2} (\langle (x, y), (u, v) \rangle + \langle (u, v), (x, y) \rangle) = \langle x, u \rangle_{\mathbb{R}} + \langle y, v \rangle_{\mathbb{R}}$$

With respect to $\langle \cdot, \cdot \rangle_{\mathbb{C}}$, $H \times H$ is $\mathbb{C}$ -vector space of dimension $2n$. With respect to $\langle \cdot, \cdot \rangle_{\mathbb{R}}$, $H \times H$ is $\mathbb{R}$ -vector space of dimension $4n$. On $H_{\mathbb{R}}$ the two inner products coincide. Because of this we shall simply denote $\langle \xi, \eta \rangle_{H_{\mathbb{R}}}$ or $\langle \xi, \eta \rangle$ the scalar product on $H \times H$ whenever $\xi, \eta \in H_{\mathbb{R}}$. Furthermore, $\mathbf{j}$ is norm preserving $\|x\|_H = \|\mathbf{j}(x)\|_{H_{\mathbb{R}}}$ and $\langle x, y \rangle_{\mathbb{R}} = \langle \mathbf{j}(x), \mathbf{j}(y) \rangle_{H_{\mathbb{R}}}$ for all $x, y \in H$. The orthogonal complement of $H_{\mathbb{R}}$ in $H \times H$ with respect to $\langle \cdot, \cdot \rangle_{\mathbb{R}}$ is given by

$$(2.22) \quad H_{\mathbb{R}}^{\perp} = iH_{\mathbb{R}} = \left\{ \left(\frac{i}{2}(x + c(x)), \frac{1}{2}(x - c(x)) \right), x \in H \right\}$$

The orthogonal projection onto $H_{\mathbb{R}}$ with respect to the real structure is given by

$$(2.23) \quad P_{\mathbb{R}} : H \times H \rightarrow H_{\mathbb{R}} \subset H \times H, \quad P_{\mathbb{R}}(u, v) = \left(\frac{1}{2}(u + c(u)), \frac{1}{2}(v + c(v)) \right)$$

The projection onto the orthogonal complement $H_{\mathbb{R}}^{\perp}$ is given by

$$(2.24) \quad P_{\mathbb{R}}^{\perp} = 1 - P_{\mathbb{R}}, \quad P_{\mathbb{R}}^{\perp}(u, v) = \left(\frac{1}{2}(u - c(u)), \frac{1}{2}(v - c(v)) \right) = i \left(\frac{1}{2i}(u - c(u)), \frac{1}{2i}(v - c(v)) \right)$$

Note:

$$(2.25) \quad P_{\mathbb{R}}^{\perp}(u, v) = -iP(iu, iv)$$

Fix $a, b \in H$ and define the linear operator on H ,

$$(2.26) \quad T_{a,b} : H \rightarrow H, \quad T_{a,b}(x) = \langle x, a \rangle b$$

Associate the following $\mathbb{R}$ -linear operator on $H \times H$

$$(2.27) \quad \tilde{T}_{a,b} : H \times H \rightarrow H_{\mathbb{R}} \subset H \times H, \quad \tilde{T}_{a,b}(u) = \langle u, \mathbf{j}(a) \rangle_{\mathbb{R}} \mathbf{j}(b) + \langle u, \mathbf{j}(ia) \rangle_{\mathbb{R}} \mathbf{j}(ib)$$

Note $\langle \mathbf{j}(b), \mathbf{j}(ib) \rangle_{\mathbb{R}} = 0$ and $H_{\mathbb{R}}$ is invariant under the action of $T_{a,b}$. Direct computations show the following diagram is commutative:

$$(2.28) \quad \begin{array}{ccccc} T_{a,b} & : & H & \xrightarrow{T_{a,b}} & H \\ & & \mathbf{j} \downarrow & & \downarrow \mathbf{j} \\ \tilde{T}_{a,b} & : & H_{\mathbb{R}} & \xrightarrow{\tilde{T}_{a,b}} & H_{\mathbb{R}} \end{array}$$

Similarly each symmetric operator in $Sym(H)$ gets mapped into a symmetric operator in $Sym(H_{\mathbb{R}})$ of double rank. Denote by τ this mapping, $\tau : Sym(H) \rightarrow Sym(H_{\mathbb{R}})$ that makes the following diagram commutative:

$$(2.29) \quad \begin{array}{ccc} H & \xrightarrow{T} & H \\ \mathbf{j} \downarrow & & \downarrow \mathbf{j} \\ H_{\mathbb{R}} & \xrightarrow{\tau(T)} & H_{\mathbb{R}} \end{array}$$

If desired, $\tau(T)$ can be extended to $H \times H$ using the $\mathbb{R}$ -linear scalar product $\langle \cdot, \cdot \rangle_{\mathbb{R}}$ on $H \times H$. τ is $\mathbb{R}$ -linear but not $\mathbb{C}$ -linear. In particular:

$$(2.30) \quad T = \llbracket x, x \rrbracket \in \mathcal{S}^{1,0} \implies \tau(T) = \llbracket \mathbf{j}(x), \mathbf{j}(x) \rrbracket + \llbracket \mathbf{j}(ix), \mathbf{j}(ix) \rrbracket \in \mathcal{S}^{2,0}(H_{\mathbb{R}})$$

$$(2.31) \quad T = \llbracket x, y \rrbracket \in \mathcal{S}^{1,1} \implies \tau(T) = \llbracket \mathbf{j}(x), \mathbf{j}(y) \rrbracket + \llbracket \mathbf{j}(ix), \mathbf{j}(iy) \rrbracket \in \mathcal{S}^{2,2}(H_{\mathbb{R}})$$

In general an operator in $\mathcal{S}^{p,q}(H)$ gets mapped into an operator in $\mathcal{S}^{2p,2q}(H_{\mathbb{R}})$. Note the map τ preserves scalar products between Sym operator up to a multiplicative constant:

$$(2.32) \quad \langle T, S \rangle_{B(H)} = 2 \langle \tau(T), \tau(S) \rangle_{B(H_{\mathbb{R}})}$$

Thus τ is a monomorphism (i.e. linear injective morphism) between $\mathbb{R}$ -vector spaces $Sym(H)$ and $Sym(H_{\mathbb{R}})$. We shall discuss in more details the linear map τ in subsection 3.2.

Note, in the case $H = \mathbb{C}^n$, $c(z) = (\bar{z}_1, \dots, \bar{z}_n)^T$, and $\langle v, w \rangle = v^T c(w)$, then $H_{\mathbb{R}} = \{(real(v), imag(v))\}, v \in \mathbb{C}^n\} = \mathbb{R}^{2n}$, and the $\mathbb{R}$ -linear map J has the block form

$$J = \begin{bmatrix} 0 & -I \\ I & 0 \end{bmatrix}$$

where I is the identity matrix of size n .

We return to the frame set $\mathbb{F} = \{f_1, \dots, f_m\}$. We let

$$(2.33) \quad \Phi_k = \tau(\llbracket f_k, f_k \rrbracket) = \llbracket \varphi_k, \varphi_k \rrbracket + \llbracket J\varphi_k, J\varphi_k \rrbracket$$

denote the image of $F_k = \llbracket f_k, f_k \rrbracket$ under τ , where $\varphi_k = \mathbf{j}(f_k)$ an element of $H_{\mathbb{R}}$. The last notation we introduce here is the following map on $H_{\mathbb{R}}$:

$$(2.34) \quad I : H_{\mathbb{R}} \rightarrow \text{Sym}(H_{\mathbb{R}}), \quad I(\xi) = \sum_{k=1}^m \Phi_k \llbracket \xi, \xi \rrbracket \Phi_k$$

As we will see later $I(\xi)$ is related to the Fisher information matrix for the measurement model (2.3), and it plays a key role in obtaining a necessary and sufficient condition of injectivity for the nonlinear map α . Since

$$\Phi_k \llbracket \xi, \xi \rrbracket \Phi_k = \llbracket \Phi_k \xi, \Phi_k \xi \rrbracket$$

$I(\xi)$ has the following more explicit form:

$$(2.35) \quad I(\xi) = \sum_{k=1}^m \llbracket v_k, v_k \rrbracket, \quad v_k = \Phi_k \xi = \langle \xi, \varphi_k \rangle_{\mathbb{R}} \varphi_k + \langle \xi, J\varphi_k \rangle_{\mathbb{R}} \varphi_k.$$

2.2. Existing Results. We revise now existing results on injectivity of the nonlinear map α . A subset Z of a topological space (X, τ) is said *generic* if its open interior is dense. In the following statements, the term *generic* refers to Zarisky topology: a set $Z \subset \mathbb{K}^{n \times m} = \mathbb{K}^n \times \cdots \times \mathbb{K}^n$ is said *generic* if Z is dense in $\mathbb{K}^{n \times m}$ and its complement is a finite union of zero sets of polynomials in nm variables with coefficients in the field $\mathbb{K}$ (here $\mathbb{K} = \mathbb{R}$ or $\mathbb{K} = \mathbb{C}$).

Theorem 2.1. *In the real case when $H = \mathbb{R}^n$ the following are equivalent:*

- (1) *The nonlinear map α is injective;*
- (2) *([4] Th.2.8) For any disjoint partition of the frame set $\mathbb{F} = \mathbb{F}_1 \cup \mathbb{F}_2$, either $\mathbb{F}_1$ spans H or $\mathbb{F}_2$ spans H .*
- (3) *([9], Th.2.4(2)) For any two vectors $x, y \in H$ if $x \neq 0$ and $y \neq 0$ then*

$$\sum_{k=1}^m |\langle x, f_k \rangle|^2 |\langle y, f_k \rangle|^2 > 0;$$

- (4) *([9], Th.2.4(3)) There is a positive real constant $a_0 > 0$ so that for all $x, y \in H$,*

$$(2.36) \quad \sum_{k=1}^m |\langle x, f_k \rangle|^2 |\langle y, f_k \rangle|^2 \geq a_0 \|x\|^2 \|y\|^2$$

- (5) *([9], Th.2.4(4)) There is a positive real constant $a_0 > 0$ so that for all $x \in H$,*

$$(2.37) \quad R(x) := \sum_{k=1}^m |\langle x, f_k \rangle|^2 \langle \cdot, f_k \rangle f_k \geq a_0 I$$

where the inequality is in the sense of quadratic forms.

Additionally, the following statements hold true:

- (1) *([4] Prop.2.5) If α is injective then $m \geq 2n - 1$;*
- (2) *([4] Prop.2.5) If $m \leq 2n - 2$ then φ cannot be injective;*

- (3) ([4] Cor.2.7(1)) If $m = 2n - 1$ then α is injective if and only if $\mathbb{F}$ is full spark;
- (4) ([4] Cor.2.6) If $m \geq 2n - 1$ and $\mathbb{F}$ is full spark then the map α is injective;
- (5) ([4] Th.2.2) If $m \geq 2n - 1$ then for a generic frame $\mathbb{F}$ the map α is injective.

In the complex case the following results are known:

Theorem 2.2. *In the complex case when $H = \mathbb{C}^n$ the following statements hold true:*

- (1) ([4] Th.3.3) If $m \geq 4n - 2$ then for a generic frame $\mathbb{F}$ the map α is injective;
- (2) ([14]) For any positive integer n there is a frame of $m = 4n - 4$ elements so that the nonlinear map α is injective;
- (3) ([38] Corollary 4) If α is injective then

$$(2.38) \quad m \geq 4n - 2 - 2\beta + \begin{cases} 2 & \text{if } n \text{ odd and } \beta = 3 \bmod 4 \\ 1 & \text{if } n \text{ odd and } \beta = 2 \bmod 4 \\ 0 & \text{otherwise} \end{cases}$$

where $\beta = \beta(n)$ denotes the number of 1's in the binary expansion of $n - 1$.

- (4) The following are equivalent:
 - (a) The map α is injective;
 - (b) ([38] Prop. 2)

$$(2.39) \quad \ker(\mathcal{A}) \cap (\mathcal{S}^{1,0} - \mathcal{S}^{1,0}) = \{0\}$$

- (c) ([11], Theorem 4) $\dim S(u) \geq 2n - 1$ for every $u \in \mathbb{C}^n \setminus \{0\}$;
- (d) ([11], Theorem 4) $S(u) = \text{span}_{\mathbb{R}}(iu)^{\perp}$ for every $u \in \mathbb{C}^n \setminus \{0\}$.

In the last two conditions, $S(u) = \text{span}_{\mathbb{R}}\{f_k f_k^* u\}_{1 \leq k \leq m}$, where $f_k f_k^* u$ is seen as a $2n$ vector in $\mathbb{R}^{2n}$.

3. NEW ANALYSIS RESULTS

This section contains our injectivity results of the nonlinear map α as well as an in-depth analysis of the spaces $\mathcal{S}^{p,q}$.

Theorem 3.1. *Let H be a $\mathbb{C}$ -vector space of dimension n , with scalar product $\langle, \rangle$ and conjugation $c : H \rightarrow H$. The following are equivalent:*

- (1) The nonlinear map $\alpha : \hat{H} \rightarrow \mathbb{R}^m$, $(\alpha(x))_k = |\langle x, f_k \rangle|^2$ is injective.
- (2) There is a positive constant $a_0 > 0$ so that for every $u, v \in H$

$$(3.40) \quad \sum_{k=1}^m |\langle F_k, \llbracket u, v \rrbracket \rangle|^2 \geq a_0 \|\llbracket u, v \rrbracket\|_1^2.$$

where $F_k = \llbracket f_k, f_k \rrbracket$. Explicitly, this means:

$$(3.41) \quad \sum_{k=1}^m (\text{real}(\langle u, f_k \rangle \langle f_k, v \rangle))^2 \geq a_0 [\|u\|^2 \|v\|^2 - (\text{imag}(\langle u, v \rangle))^2].$$

- (3) For any $\xi \in H_R$, $\xi \neq 0$, $\text{rank}(I(\xi)) = 2n - 1$.

(4) There is $a_0 > 0$ so that for all $\xi \in H_{\mathbb{R}}$, $\xi \neq 0$,

$$(3.42) \quad I(\xi) \geq a_0 \|\xi\|^2 P_{J\xi}^\perp$$

where the inequality holds in the sense of quadratic forms in $H_{\mathbb{R}}$, and

$$(3.43) \quad P_{J\xi}^\perp = 1 - \frac{1}{\|\xi\|^2} \llbracket J\xi, J\xi \rrbracket$$

is the orthogonal projection in $H_{\mathbb{R}}$ onto the orthogonal complement of $J\xi$ in $H_{\mathbb{R}}$.

Remark 3.2. The two constants a_0 in (2) and (4) can be chosen to be equal, hence the same notation. We see in the next section, this common constant is related to robustness and stability of any reconstruction scheme.

Remark 3.3. The proof of (3) $\Leftrightarrow$ (4) shows that the optimal bound a_0 is given by

$$(3.44) \quad a_0^{\text{opt}} = \min_{\xi \in H_{\mathbb{R}}, \|\xi\|=1} a_{n-1}(I(\xi))$$

where $a_{n-1}(I(\xi))$ denotes the next to the smallest eigenvalue of $I(\xi)$.

Remark 3.4. The choice of the nuclear norm and the square in (3.40) is somewhat arbitrary. For any $p, q \geq 1$ (including infinity), there is a constant $a_{p,q} > 0$ so that

$$(3.45) \quad \sum_{k=1}^m |\langle F_k, \llbracket u, v \rrbracket \rangle|^p \geq a_{p,q} \|\llbracket u, v \rrbracket\|_q^p$$

An interesting corollary is obtained in the case when α is restricted to a subspace of H . Specifically there is sometime the case the underlying signal is real. The canonical description of such a condition is to be conjugation invariant. Let H' denote this set:

$$(3.46) \quad H' = \{x \in H ; c(x) = x\}$$

Note H' is a $\mathbb{R}$ -linear space, but it is not a $\mathbb{C}$ -linear space. Since x is restricted to H' it follows the equivalence class in H' of x is given by $\hat{x} = \{x, -x\}$. Consequently the appropriate quotient space is given by $\widehat{H'} = \{\{x, -x\}, x \in H_{\mathbb{R}}\}$. Let $\pi_1 : H \times H \rightarrow H$ and $\pi_2 : H \times H \rightarrow H$ be the canonical projections onto factors: $\pi_1((u, v)) = u$ and $\pi_2((u, v)) = v$. Then it is immediate to check that H' admits the following equivalent descriptions:

$$(3.47) \quad H' = \{x \in H ; \pi_2(\mathbf{j}(x)) = 0\} = \mathbf{j}^{-1}(\{H_{\mathbb{R}} \cap (H \times \{0\})\})$$

Note in H' , $\langle u, v \rangle$ is always real for all $u, v \in H'$ since $\langle u, v \rangle = \langle c(v), c(u) \rangle = \langle v, u \rangle$. Let $\mathbb{F} = \{f_1, \dots, f_m\}$ be the frame set in H . Let

$$(3.48) \quad g_k = \pi_1(\mathbf{j}(f_k)) , \quad h_k = \pi_2(\mathbf{j}(f_k))$$

with $1 \leq k \leq m$. Note $f_k = \mathbf{j}^{-1}((g_k, h_k)) = g_k + ih_k$ and $g_k, h_k \in H'$. Set:

$$(3.49) \quad \Phi'_k = \llbracket g_k, g_k \rrbracket + \llbracket h_k, h_k \rrbracket \in \mathcal{S}^{2,0}(H') \subset \mathcal{S}^{2,0}(H) , \quad 1 \leq k \leq m$$

Note:

$$(3.50) \quad \langle F_k, \llbracket x, x \rrbracket \rangle = |\langle f_k, x \rangle|^2 = \langle \Phi'_k x, x \rangle = \langle \Phi'_k, \llbracket x, x \rrbracket \rangle$$

$$(3.51) \quad \langle F_k, \llbracket u, v \rrbracket \rangle = \text{real}(\langle u, f_k \rangle \langle f_k, v \rangle) = \langle \Phi'_k u, v \rangle = \langle \Phi'_k, \llbracket u, v \rrbracket \rangle$$

for all $x, u, v \in H'$. Thus the linear map $\mathcal{A}$ restricted to $\text{Sym}(H')$ can be thought of as taking inner products with a family of rank-2 nonnegative operators. We have

Corollary 3.5. *Let H be a n -dimensional complex Hilbert space with scalar product $\langle, \rangle$ and conjugation $c : H \rightarrow H$. Let $H' = \{x \in H ; c(x) = x\}$ be the c -invariant set. The following are equivalent:*

- (1) *The restriction to H' of the nonlinear map $\alpha|_{H'}$ is injective on $\widehat{H'}$.*
- (2) *There is a constant $a_0 > 0$ so that*

$$(3.52) \quad \sum_{k=1}^m |\langle \Phi_k u, v \rangle|^2 \geq a_0 \|u\|^2 \|v\|^2$$

- (3) *For all $u \neq 0$,*

$$(3.53) \quad \dim_{\mathbb{R}} \text{span}_{\mathbb{R}} \{ \langle u, g_k \rangle g_k + \langle u, h_k \rangle h_k ; 1 \leq k \leq m \} = n$$

- (4) *For any $x \neq 0$,*

$$(3.54) \quad \text{rank} \left(\sum_{k=1}^m \Phi'_k \llbracket x, x \rrbracket \Phi'_k \right) = n$$

- (5) *There is a constant $a_0 > 0$ so that*

$$(3.55) \quad I'(x) = \sum_{k=1}^m \Phi'_k \llbracket x, x \rrbracket \Phi'_k \geq a_0 \|x\|^2 1_{H'}$$

in the sense of quadratic forms on H' , where $1_{H'}$ is the identity map on H' .

3.1. Analysis of sets $\mathcal{S}^{p,q}$. In addition to (2.11) the sets $\mathcal{S}^{p,q}$ introduced in (2.7) have the following properties summarized in the following lemma:

Lemma 3.6. (1) *For any $p_1 \leq p_2$ and $q_1 \leq q_2$, $\mathcal{S}^{p_1, q_1} \subset \mathcal{S}^{p_2, q_2}$*

- (2) *For any nonnegative integers p, q the following disjoint decomposition holds true*

$$(3.56) \quad \mathcal{S}^{p,q} = \bigcup_{r=0}^p \bigcup_{s=0}^q \mathring{\mathcal{S}}^{r,s}$$

where by convention $\mathring{\mathcal{S}}^{0,0} = \mathcal{S}^{0,0} = \{0\}$, and $\mathring{\mathcal{S}}^{p,q} = \emptyset$ for $p+q > n$.

- (3) *For any nonnegative integers p, q ,*

$$(3.57) \quad - \mathcal{S}^{p,q} = \mathcal{S}^{q,p}$$

(4) For any bounded $T \in B(H)$ and integers p, q ,

$$(3.58) \quad T^* \mathcal{S}^{p,q} T \subset \mathcal{S}^{p,q}$$

The inclusion becomes equality if and only if T is invertible.

(5) For any integers p, q, r, s ,

$$(3.59) \quad \mathcal{S}^{p,q} + \mathcal{S}^{r,s} = \mathcal{S}^{p,q} - \mathcal{S}^{s,r} = \mathcal{S}^{p+r,q+s}$$

Proof of Lemma 3.6

First three assertions are trivial. Assertions (4) & (5) state that the only invariant of a quadratic form over $\mathbb{C}$ is its signature. Assertion (4) can be shown using the singular value decomposition of T and the continuity of spectrum with respect to matrix entries. For (5), one can see this statement as a special instance of the Witt decomposition theorem, a much more powerful tool in the theory of quadratic forms especially in the case of vector spaces over fields of characteristics other than 2, see [46]. However for the benefit of those who prefer a more direct proof, here is a sketch of this result. Using spectral decomposition and rearranging the term, one can easily see that $\mathcal{S}^{p+r,q+s} \subset \mathcal{S}^{p,q} + \mathcal{S}^{r,s}$. For the converse inclusion, we need to show that if $T, S \in \text{Sym}(H)$ are so that T has at most p positive eigenvalues and q negative eigenvalues, and S has at most r positive eigenvalues and s negative eigenvalues, then $T + S$ has at most $p + r$ positive eigenvalues, and $q + s$ negative eigenvalues. Without loss of generality we can assume $T \in \mathring{\mathcal{S}}^{p,q}$ and $S \in \mathring{\mathcal{S}}^{r,s}$. Using spectral decompositions of T and S we obtain

$$\begin{aligned} T + S &= \left(\sum_{k=1}^p a_k A_k - \sum_{k=p+1}^{p+q} a_k A_k \right) + \left(\sum_{k=1}^r b_k B_k - \sum_{k=r+1}^{r+s} b_k B_k \right) = U - V, \\ U &= \sum_{k=1}^p a_k A_k + \sum_{k=1}^r b_k B_k \geq 0, \quad V = \sum_{k=p+1}^q a_k A_k + \sum_{k=r+1}^{r+s} b_k B_k \geq 0, \end{aligned}$$

where A_k, B_k are rank one orthogonal (spectral) projectors with $A_k A_j = 0$ and $B_l B_h = 0$ for all $k \neq j$ and $l \neq h$, and $a_k, b_k > 0$. Thus $U \in \mathcal{S}^{p+r,0}$ and $V \in \mathcal{S}^{q+s,0}$. The claim now follows by induction provided we show that for any $R \in \mathring{\mathcal{S}}^{a,b}$ and $E \in \mathcal{S}^{1,0}$ a rank one associated to vector $e \in H$ ($E = \langle \cdot, e \rangle e$), then $E + R \in \mathcal{S}^{a+1,b}$.

Indeed this last assertion is shown as follows. Let $R = \sum_{k=1}^a c_k \langle \cdot, g_k \rangle g_k - \sum_{k=1+a}^{a+b} c_k \langle \cdot, g_k \rangle g_k$ be its spectral decomposition. Two cases are treated distinctly.

Case 1: If $e \notin R(H)$ then $\Gamma' = \{g_1, \dots, g_{a+b}, e\}$ is linearly independent. Let $\Gamma = \{\gamma_k, 1 \leq k \leq a+b+1\}$ be the (unique) biorthogonal system to Γ' . Let T be an invertible operator that maps an orthonormal set $\Delta = \{\delta_1, \dots, \delta_{a+b+1}\}$ into Γ , $T\delta_k = \gamma_k$, $1 \leq k \leq a+b+1$, and maps the orthogonal space to Δ onto the orthogonal complement to Γ . Note $T^* g_k = \delta_k$, $1 \leq k \leq a+b$ and $T^* e = \delta_{a+b+1}$. Then a direct computation shows that

$$T^*(R + E)T = \sum_{k=1}^{a+b} c_k \langle \cdot, \delta_k \rangle \delta_k + \langle \cdot, \delta_{a+b+1} \rangle \delta_{a+b+1}$$

Thus the spectrum of $T^*(R+E)T$ is composed of $\{c_1, \dots, c_a, -c_{a+1}, \dots, c_{a+b}, 1\}$ which shows that $T^*(R+E)T \in \mathcal{S}^{a+1,b}$, and by (4), $R+E \in \mathcal{S}^{a+1,b}$.

Case 2: $e \in R(H)$. The rank of $R+E$ is the same as the rank of R , hence $\dim \ker(R+E) = \dim \ker(R)$. Hence $R+E \in \mathcal{S}^{a',b'}$ with $a'+b' = a+b$. Now by the continuity of spectrum with respect to small perturbations, it follows that, for a small perturbation $e \mapsto e' \notin R(H)$, $R+E' \in \mathcal{S}^{a'',b''}$ with $a'' \geq a', b'' \geq b'$. But the proof of case 1 shows $\mathcal{S}^{a'',b''} \subset \mathcal{S}^{a+1,b}$. Hence $b \geq b'' \geq b' \geq b$, and $a+1 \geq a'' \geq a' \geq a$. Thus $R+E \in \mathcal{S}^{a+1,b}$. Q.E.D. $\square$.

Next we analyze the space $\mathcal{S}^{1,1}$ in more detail. The special factorization we obtain here can be extended to other classes $\mathcal{S}^{p,q}$ but we do not plan to do so here. First set the following matrices:

$$(3.60) \quad H = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad D = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \quad V = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}.$$

In the next result we use a generalized unitary group $U(1,1;H)$. Recall its definition.

Definition The groups $U(1,1)$ and $U(1,1;H)$ are defined by

$$(3.61) \quad U(1,1) = \{A \in \mathbb{C}^{2 \times 2}, A^*DA = D\}$$

$$(3.62) \quad U(1,1;H) = \{A \in \mathbb{C}^{2 \times 2}, A^*HA = H\}$$

These groups been studied extensively in literature. See for instance [56], section 10.4. In particular the two groups above are unitarily equivalent to each other, and the matrix V provides such an equivalence:

$$(3.63) \quad A \in U(1,1;H) \Leftrightarrow B = VAV^* \in U(1,1)$$

The quadratic form $\omega(z_1, z_2) = |z_1|^2 - |z_2|^2$ is invariant under the action of $U(1,1)$, whereas $\phi(z_1, z_2) = \bar{z}_1 z_2 + z_1 \bar{z}_2$ is invariant under the action of $U(1,1;H)$.

Lemma 3.7. (1) $\mathcal{S}^{1,1} = \mathcal{S}^{1,0} - \mathcal{S}^{1,0} = \mathcal{S}^{1,0} + \mathcal{S}^{0,1}$.

(2) For any $T \in \mathcal{S}^{1,1}$ there are $u, v \in H$ so that

$$(3.64) \quad T = \frac{1}{2} (\langle \cdot, u \rangle v + \langle \cdot, v \rangle u) = \llbracket u, v \rrbracket$$

where the symmetric outer product $\llbracket u, v \rrbracket$ has been defined in (2.5).

If $T = a_1 \llbracket e_1, e_1 \rrbracket - a_2 \llbracket e_2, e_2 \rrbracket$ with $a_1, a_2 \geq 0$ and $\langle e_k, e_j \rangle = \delta_{k,j}$ is its spectral factorization then

$$(3.65) \quad u_0 = \sqrt{a_1}e_1 + \sqrt{a_2}e_2, \quad v_0 = \sqrt{a_1}e_1 - \sqrt{a_2}e_2$$

provides a particular factorization in (3.64). Conversely, for any $u, v \in H$, the operator $T = \llbracket u, v \rrbracket$ is in $\mathcal{S}^{1,1}$.

(3) Let $T = \llbracket u_0, v_0 \rrbracket \in \mathcal{S}^{1,1}$. Then any pair (u, v) of vectors in H so that $T = \llbracket u, v \rrbracket$ is given by

$$(3.66) \quad u = a_{11}u_0 + a_{12}v_0, \quad v = a_{21}u_0 + a_{22}v_0$$

for some matrix $A = (a_{k,l})_{1 \leq k,l \leq 2}$ with $A \in U(1, 1; H)$. Conversely, for any matrix $A \in U(1, 1; H)$, $\llbracket u, v \rrbracket = \llbracket u_0, v_0 \rrbracket$ where (u, v) are given by (3.66).

(4) Let $T = \llbracket u, v \rrbracket$. Then traces and spectrum $Sp(T) = \{a_+, a_-\}$ are given by

$$(3.67) \quad tr\{T\} = real(\langle u, v \rangle) = \langle u, v \rangle_{\mathbb{R}}$$

$$(3.68) \quad \begin{aligned} tr\{T^2\} &= \frac{1}{4}((\langle u, v \rangle)^2 + (\langle v, u \rangle)^2 + 2\|u\|^2\|v\|^2) \\ &= \frac{1}{2}(\|u\|^2\|v\|^2 + \langle u, v \rangle_{\mathbb{R}}^2 - \langle iu, v \rangle_{\mathbb{R}}^2) \end{aligned}$$

$$(3.69) \quad a_+ = \frac{1}{2} \left(\langle u, v \rangle_{\mathbb{R}} + \sqrt{\|u\|^2\|v\|^2 + \langle iu, v \rangle_{\mathbb{R}}^2} \right) \geq 0$$

$$(3.70) \quad a_- = \frac{1}{2} \left(\langle u, v \rangle_{\mathbb{R}} - \sqrt{\|u\|^2\|v\|^2 + \langle iu, v \rangle_{\mathbb{R}}^2} \right) \leq 0$$

Hence the nuclear norm of T is given by

$$(3.71) \quad \|T\|_1 = a_+ + |a_-| = \sqrt{\|u\|^2\|v\|^2 + \langle iu, v \rangle_{\mathbb{R}}^2}$$

Remark 3.8. 1. The need for studying $\mathcal{S}^{1,1}$ arose from the behavior of the IRLS algorithm described in [9]. However it quickly become apparent the space $\mathcal{S}^{1,1}$ and its factorization given by (3.64) are crucial for understanding the injectivity of the nonlinear map α especially in the light of Theorem 2.2 (3), equation (2.39).

2. The choice in (3.65) has the following two additional properties:

$$(3.72) \quad \|u_0\| = \|v_0\| = \sqrt{a_1 + a_2} = \sqrt{a_+ - a_-} = \sqrt{\|T\|_1}$$

$$(3.73) \quad \langle u_0, v_0 \rangle = a_1 - a_2 = a_+ + a_- = tr\{T\} \text{ (a real number!)}$$

where $\|T\|_1$ represents the nuclear norm of T (sum of its singular eigenvalues), and $a_1 = a_+ \geq 0$ and $a_2 = -a_- \geq 0$ are its singular eigenvalues.

Proof of Lemma 3.7

(1) is a direct application of Lemma 3.6(5).

(2) Since $\llbracket \cdot, \cdot \rrbracket$ is $\mathbb{R}$ -linear and $\llbracket e_1, e_2 \rrbracket = \llbracket e_2, e_1 \rrbracket$ we obtain

$$\llbracket u_0, v_0 \rrbracket = a_1 \llbracket e_1, e_1 \rrbracket + \sqrt{a_1 a_2} \llbracket e_1, e_2 \rrbracket - \sqrt{a_1 a_2} \llbracket e_1, e_1 \rrbracket - a_2 \llbracket e_2, e_2 \rrbracket = a_1 \llbracket e_1, e_1 \rrbracket - a_2 \llbracket e_2, e_2 \rrbracket = T$$

For the converse, note $rank(T) = dim \text{span}\{u, v\} \leq 2$. Then the claim follows once we establish (3.69) and (3.70) which we do below.

(3) A direct computation shows

$$\begin{aligned} \llbracket u, v \rrbracket &= \frac{1}{2}(\bar{a}_{11}a_{21} + \bar{a}_{21}a_{11})\langle \cdot, u_0 \rangle u_0 + \frac{1}{2}(\bar{a}_{11}a_{22} + \bar{a}_{21}a_{12})\langle \cdot, u_0 \rangle v_0 \\ &\quad + \frac{1}{2}(\bar{a}_{12}a_{21} + \bar{a}_{22}a_{11})\langle \cdot, v_0 \rangle u_0 + \frac{1}{2}(\bar{a}_{12}a_{22} + \bar{a}_{22}a_{12})\langle \cdot, v_0 \rangle v_0 \end{aligned}$$

Since $\{u_0, v_0\}$ are linearly independent, $\llbracket u, v \rrbracket = \llbracket u_0, v_0 \rrbracket$ implies

$$\bar{a}_{11}a_{21} + \bar{a}_{21}a_{11} = 0, \quad \bar{a}_{11}a_{22} + \bar{a}_{21}a_{12} = 1$$

$$\bar{a}_{12}a_{21} + \bar{a}_{22}a_{11} = 1, \quad \bar{a}_{12}a_{22} + \bar{a}_{22}a_{12} = 0$$

Which corresponds to $A^*HA = H$. Hence $A \in U(1, 1; H)$. Conversely, if $A \in U(1, 1; H)$ then above relations are satisfied which imply $\llbracket u, v \rrbracket = \llbracket u_0, v_0 \rrbracket$.

(4) The equation (3.67) comes from the definition (2.5) and the fact that $\text{tr}(\langle \cdot, u \rangle v) = \langle v, u \rangle$. For T^2 compute first

$$T^2 = \frac{1}{4} (\langle v, u \rangle \langle \cdot, u \rangle v + \|u\|^2 \langle \cdot, v \rangle v + \|v\|^2 \langle \cdot, u \rangle u + \langle u, v \rangle \langle \cdot, v \rangle u).$$

Then (3.68) follows from this relation and $\text{real}((\langle u, v \rangle)^2) = (\text{real}(\langle u, v \rangle))^2 - (\text{imag}(\langle u, v \rangle))^2$. Finally, (3.69) and (3.70) come from solving:

$$(3.74) \quad \begin{aligned} a_+ + a_- &= \text{tr}(T) \\ a_+^2 + a_-^2 &= \text{tr}(T^2) \end{aligned}$$

Q.E.D. $\square$

Topologically, $\mathcal{S}^{1,0}$ and $\mathcal{S}^{1,1}$ are not differentiable manifolds. Instead they are algebraic varieties, since are given by the zero loci of certain polynomials. We have the following result:

Lemma 3.9. (1) *The set $\mathring{\mathcal{S}}^{1,0}$ is an analytic manifold in $B(H)$ of real dimension $2n - 1$. As real manifold, its tangent space at $X = \llbracket x, x \rrbracket$ is given by*

$$(3.75) \quad T_X \mathring{\mathcal{S}}^{1,0} = \{\llbracket x, y \rrbracket, y \in H\}$$

The $\mathbb{R}$ -linear embedding $H \mapsto T_X \mathring{\mathcal{S}}^{1,0}$ given by $y \mapsto \varphi_x(y) = \llbracket x, y \rrbracket$ has null space given by $\ker \varphi_x = \{iax; a \in \mathbb{R}\}$.

(2) *The set $\mathring{\mathcal{S}}^{1,1}$ is an analytic manifold of real dimension $4n - 3$. As real manifold, its tangent space at $T = \llbracket x, y \rrbracket$ is given by*

$$(3.76) \quad T_T \mathring{\mathcal{S}}^{1,1} = \{\llbracket x, u \rrbracket + \llbracket y, v \rrbracket, u, v \in H\}$$

The $\mathbb{R}$ -linear embedding $H \times H \mapsto T_T \mathring{\mathcal{S}}^{1,1}$ given by $(x, y) \mapsto \varphi_{x,y}(u, v) = \llbracket x, u \rrbracket + \llbracket y, v \rrbracket$ has null space given by $\ker \varphi_{x,y} = \{a(ix, 0) + b(0, iy) + c(y, -x), a, b, c \in \mathbb{R}\}$.

Proof of Lemma 3.9

Let $c_1, \dots, c_n : \text{Sym}(H) \rightarrow \mathbb{R}$ be the coefficients of the characteristic polynomial:

$$\det(sI - T) = s^n + c_1(T)s^{n-1} + c_2(T)s^{n-2} + \dots + c_n(T)$$

Thus $c_1(T) = -\text{tr}(T)$ and $c_n(T) = (-1)^n \det(T)$.

(1) The manifold structure can be shown as follows. First note that

$$S^+ = \{S \in \text{Sym}(H); c_1(T) = -\text{tr}(S) < 0\}$$

is an open subset of $Sym(H)$ and therefore a manifold of same real dimension as $Sym(H)$ (which is $2n$). Next note

$$\mathring{\mathcal{S}}^{1,0} = c_2^{-1}(0) \cap \cdots \cap c_n^{-1}(0) \cap S^+$$

Hence $\mathring{\mathcal{S}}^{1,0}$ is an algebraic variety. We obtain further it is an analytic manifold by showing its tangent space has same dimension at all points. Let $X = \llbracket x_0, x_0 \rrbracket \in \mathring{\mathcal{S}}^{1,0}$. We consider the set of all differentiable curves $\gamma : I \rightarrow \mathring{\mathcal{S}}^{1,0}$, $\gamma(0) = X$, $0 \in I \subset \mathbb{R}$ open, passing through X . Then the tangent space at X is given by

$$T_X \mathring{\mathcal{S}}^{1,0} = \left\{ \frac{d}{dt} \gamma(t) \Big|_{t=0} \right\}$$

For each such curve, $\gamma : I \rightarrow \mathring{\mathcal{S}}^{1,0}$ there is a unique curve $x : I \rightarrow H$ such that $\gamma(t) = \llbracket x(t), x(t) \rrbracket$ with $x(0) = x_0$. In fact, locally,

$$x(t) = \frac{1}{\sqrt{\langle \gamma(t)(x_0), x_0 \rangle}} \gamma(t)(x_0)$$

which shows $x(t)$ is differentiable. Then a direct computation shows

$$\frac{d}{dt} \gamma(t) \Big|_{t=0} = \llbracket \dot{x}(0), x(0) \rrbracket + \llbracket x(0), \dot{x}(0) \rrbracket = 2 \llbracket x_0, \dot{x}(0) \rrbracket$$

Since $\dot{x}(0)$ can be chosen arbitrarily in H , it follows the tangent space to $\mathring{\mathcal{S}}^{1,0}$ at X is given by (3.75). The real dimension of the $\mathbb{R}$ -vector space $T_X \mathring{\mathcal{S}}^{1,0}$ is $2n - 1$ once we notice the kernel of the $\mathbb{R}$ -linear map φ_x is one dimensional and given by the span of ix .

(2) The algebraic manifold structure is given by the intersection

$$\mathring{\mathcal{S}}^{1,1} = c_3^{-1}(0) \cap \cdots \cap c_n^{-1}(0) \cap S^{--}$$

where

$$S^{--} = \{S \in Sym(H) ; c_2(T) < 0\}$$

The analytic manifold structure follows once we establish the tangent space at every point is given by (3.76) and has same dimension $4n - 3$. Fix $T \in \mathring{\mathcal{S}}^{1,1}$, $T = \llbracket x_0, y_0 \rrbracket$ with $\{x_0, y_0\}$ linearly independent, and let $\{\gamma : I \rightarrow \mathring{\mathcal{S}}^{1,1}, 0 \in I \subset \mathbb{R} \text{ open}, \gamma(0) = T\}$ be the set of differentiable curves passing through T . By Lemma 3.6 we know $\gamma(t) = \llbracket x(t), y(t) \rrbracket$ for some $x : I \rightarrow H$ and $y : I \rightarrow H$. However these functions are not unique. However we can choose them to be given by the spectral factorization of $\gamma(t)$ via (3.65). Furthermore a direct application of functional calculus (see [53]) shows that $T \rightarrow P_1 \in \mathring{\mathcal{S}}^{1,0}$ and $T \mapsto P_2 \in \mathring{\mathcal{S}}^{1,0}$ are analytic, where $T = P_1 - P_2$ is its spectral decomposition with $P_1 P_2 = 0$. Hence $t \in I \mapsto P_1(t)$ and $t \in I \mapsto P_2(t)$ are differentiable. Then the component functions $x(t)$ and $y(t)$ now uniquely defined by:

$$(3.77) \quad x(t) = \frac{1}{\sqrt{\langle P_1(t)(x(0)), x(0) \rangle}} P_1(t)(x(0)) , \quad y(t) = \frac{1}{\sqrt{\langle P_2(t)(y(0)), y(0) \rangle}} P_2(t)(y(0))$$

are differentiable. The derivative at $t = 0$ is given by

$$\frac{d}{dt}\gamma(t)|_{t=0} = 2\llbracket x(0), \dot{y}(0) \rrbracket + 2\llbracket y(0), \dot{x}(0) \rrbracket$$

And since $(x(0), y(0))$ and (x_0, y_0) are related by a $U(1, 1; H)$ matrix, and the fact that $(\dot{x}(0), \dot{y}(0))$ is arbitrary in $H \times H$, we obtain the tangent space is given by (3.76). A direct computation shows that $(ix, 0)$, $(0, iy)$ and $(y, -x)$ are independent vectors in the null space of $\varphi_{x,y}$ which concludes the proof. $\square$

Remark 3.10. Compared to the complex projective manifold $\mathbb{C}P^n$, $\mathring{\mathcal{S}}^{1,0}$ is diffeomorphic to $\mathbb{C}P^n \times \mathbb{R}^+$. The extra $\mathbb{R}^+$ component comes from the fact that rank-1 operators in $\mathring{\mathcal{S}}^{1,0}$ have arbitrary trace. This explains the real dimension of $\mathring{\mathcal{S}}^{1,0}$: $2\dim_{\mathbb{R}} \mathbb{C}P^n + 1 = 2(n-1) + 1 = 2n-1$. On the other hand, using spectral factorization, $\mathring{\mathcal{S}}^{1,1}$ has real dimension given by $\dim_{\mathbb{R}} \mathring{\mathcal{S}}^{1,0} + \dim_{\mathbb{R}} \mathring{\mathcal{S}}^{1,0} - 1 = 2(2n-1) - 1 = 4n-3$. The -1 term comes from the orthogonality of the two eigenvectors of an operator in $\mathring{\mathcal{S}}^{1,1}$.

Remark 3.11. As suggested by Bernhard Bodmann [13], it can be shown the subset of projections inside $\mathring{\mathcal{S}}^{1,0}$ is in fact a Kähler manifold (diffeomorphic to $\mathbb{C}P^n$). A similar conclusion holds for $\mathring{\mathcal{S}}^{1,1}$ since the set of projections inside $\mathring{\mathcal{S}}^{1,1}$ is the same to the set of projections inside $\mathring{\mathcal{S}}^{1,0}$. However none of these two manifolds, $\mathring{\mathcal{S}}^{1,0}$ and $\mathring{\mathcal{S}}^{1,1}$, is Kähler: they have odd real dimension, hence cannot admit a symplectic structure.

3.2. Analysis of the linear map τ . We introduced earlier the $\mathbb{R}$ -linear map τ that maps $Sym(H)$ operators into $Sym(H_{\mathbb{R}})$ operators, using the real linear structure on these spaces. In order for the diagram (2.29) to be commutative, the map $\tau(T)$ is given explicitly by

$$(3.78) \quad \tau(T)(\xi) = \mathbf{j}(T(\mathbf{j}^{-1}(\xi))) , \quad \xi \in H_{\mathbb{R}}$$

The followin lemma summarizes the basic properties of the map τ .

Lemma 3.12. (1) Let P be an orthogonal projection of rank k in H . Then $\tau(P)$ is an orthogonal projection of rank $2k$ in $H_{\mathbb{R}}$. Furthermore if $\{e_1, \dots, e_k\}$ is an orthonormal basis in the range of P , then $\{\mathbf{j}(e_1), \dots, \mathbf{j}(e_k), J\mathbf{j}(e_1), \dots, J\mathbf{j}(e_k)\}$ is an orthonormal basis in the range of $\tau(P)$.

(2) If $T \in Sym(H)$ has spectrum $(a_1, a_2, \dots, a_n)$ then $\tau(T)$ in $B(H_{\mathbb{R}})$ has spectrum $(a_1, a_1, a_2, a_2, \dots, a_n, a_n)$.

(3) For any two operators $T, S \in Sym(H)$, $\tau(T), \tau(S) \in Sym(H_{\mathbb{R}})$ and

$$(3.79) \quad tr\{\tau(T)\tau(S)\} = \langle \tau(T), \tau(S) \rangle_{B(H_{\mathbb{R}})} = 2\langle T, S \rangle_{B(H)} = 2tr\{TS\}$$

(4) Let $1 \leq p \leq \infty$. The p -norms of a symmetric operator $T \in Sym(H)$ and $\tau(T) \in Sym(H_{\mathbb{R}})$, are related by

$$(3.80) \quad \|\tau(T)\|_p = 2^{1/p} \|T\|_p , \quad \text{if } p < \infty$$

$$(3.81) \quad \|T\| = \|\tau(T)\|_{\infty} = \|T\|_{\infty} = \|T\|$$

Proof of Lemma 3.12

(1) First we prove the statement for rank-1 projectors. This comes from directly checking equation (2.30). Thus $P = \llbracket e, e \rrbracket$ gets mapped into $\tau(P) = \llbracket \xi, \xi \rrbracket + \llbracket J\xi, J\xi \rrbracket$, where $\xi = \mathbf{j}(e)$. Next if $\{e_1, \dots, e_k\}$ is an orthonormal basis for the range of P then

$$P = \sum_{l=1}^k \llbracket e_l, e_l \rrbracket$$

By $\mathbb{R}$ -linearity, $\tau(P)$ has the form

$$\tau(P) = \sum_{l=1}^k \llbracket \xi_l, \xi_l \rrbracket + \llbracket J\xi_l, J\xi_l \rrbracket$$

where $\xi_l = \mathbf{j}(e_l)$, $1 \leq l \leq k$. Note

$$\langle J\xi_l, J\xi_s \rangle = \langle \xi_l, \xi_s \rangle = \langle e_l, e_s \rangle_{\mathbb{R}} = \delta_{l,s}, \quad \langle J\xi_l, \xi_s \rangle = \langle ie_l, e_s \rangle_{\mathbb{R}} = 0$$

Hence $\{\xi_1, \dots, \xi_k, J\xi_1, \dots, J\xi_k\}$ is an orthonormal set and since it is spanning the range of $\tau(P)$ it follows it is an orthonormal basis in the range of $\tau(P)$. Hence $\tau(P)$ is an orthonormal projection on $H_{\mathbb{R}}$ of rank $2k$.

(2) Follows by using the spectral factorization of T ,

$$T = \sum_{l=1}^s a_l P_l \mapsto \tau(T) = \sum_{l=1}^s a_l \tau(P_l)$$

Then $\tau(P_k)\tau(P_l) = 0$ for $k \neq l$. Hence the equation for $\tau(T)$ above represents its spectral factorization. the conclusion follows.

(3),(4) Follow from (2). □

3.3. Proof of Theorem 3.1 and its Corollary 3.5. (1) $\Leftrightarrow$ (2). According to Theorem 2.2(4), nonlinear map α is injective iff $\ker(\mathcal{A}) \cap (\mathcal{S}^{1,0} - \mathcal{S}^{1,0}) = \{0\}$. But using Lemma 3.7(1) and (2) we get equivalently that α is injective iff for all $u, v \in H$ with $\llbracket u, v \rrbracket \neq 0$,

$$\mathcal{A}(\llbracket u, v \rrbracket) \neq 0$$

Equivalently this means

$$\sum_{k=1}^m |\langle F_k, \llbracket u, v \rrbracket \rangle|^2 > 0$$

Consider now the unit ball in $\mathcal{S}^{1,1}$ with respect to the nuclear norm, say $S_1^{1,1}$. This set is compact in $Sym(H)$. Then let

$$(3.82) \quad a_0 = \min_{T \in \mathcal{S}^{1,1}, \|T\|_1=1} \sum_{k=1}^m |\langle F_k, T \rangle|^2$$

By homogeneity we obtain (3.40). Then

$$\langle F_k, \llbracket u, v \rrbracket \rangle = \frac{1}{2}(\langle u, f_k \rangle \langle f_k, v \rangle + \langle v, f_k \rangle \langle f_k, u \rangle) = \text{real}(\langle u, f_k \rangle \langle f_k, v \rangle)$$

and by Lemma 3.7(4),

$$\|\llbracket u, v \rrbracket\|_1^2 = \|u\|^2 \|v\|^2 - \langle iu, v \rangle_{\mathbb{R}}^2 = \|u\|^2 \|v\|^2 - (\text{imag}(\langle u, v \rangle))^2$$

Putting together all previous derivations we obtain the equivalence (1) $\Leftrightarrow$ (2).

(2) $\Leftrightarrow$ (4). Using Lemma 3.9(3) we obtain (3.40) is equivalent to

$$(3.83) \quad \sum_{k=1}^m |\langle \tau(F_k), \tau(\llbracket u, v \rrbracket) \rangle|^2 \geq 4a_0 [\|u\|^2 \|v\|^2 - (\text{real}(\langle iu, v \rangle))^2]$$

Now (2.30) and (2.31) imply

$$\tau(F_k) = \llbracket \varphi_k, \varphi_k \rrbracket + \llbracket J\varphi_k, J\varphi_k \rrbracket = \llbracket \varphi_k, \varphi_k \rrbracket + J\llbracket \varphi_k, \varphi_k \rrbracket J^*$$

$$\tau(\llbracket u, v \rrbracket) = \llbracket \xi, \eta \rrbracket + \llbracket J\xi, J\eta \rrbracket = \llbracket \xi, \eta \rrbracket + J\llbracket \xi, \eta \rrbracket J^*$$

where $\varphi = \mathbf{j}(f_k)$ and $\xi = \mathbf{j}(u)$, $\eta = \mathbf{j}(v)$. A direct computation using $J^* = -J$ shows that

$$\langle \tau(F_k), \llbracket J\xi, J\eta \rrbracket \rangle = \langle \tau(F_k), \llbracket \xi, \eta \rrbracket \rangle$$

Thus

$$\langle \tau(F_k), \tau(\llbracket u, v \rrbracket) \rangle = 2\langle \llbracket \varphi_k, \varphi_k \rrbracket + \llbracket J\varphi_k, J\varphi_k \rrbracket, \llbracket \xi, \eta \rrbracket \rangle = 2[\langle \xi, \varphi_k \rangle \langle \varphi_k, \eta \rangle + \langle \xi, J\varphi_k \rangle \langle J\varphi_k, \eta \rangle]$$

With the equation (2.35) we obtain

$$\langle \tau(F_k), \tau(\llbracket u, v \rrbracket) \rangle = 2\langle v_k, \eta \rangle \Rightarrow \sum_{k=1}^m |\langle \tau(F_k), \tau(\llbracket u, v \rrbracket) \rangle|^2 = 4\langle I(\xi)\eta, \eta \rangle$$

Now the right-hand side of (3.83) is processed as follows. Note $\|u\| = \|\xi\|$, $\|v\| = \|\eta\|$, and

$$\text{real}(\langle iu, v \rangle) = \langle iu, v \rangle_{\mathbb{R}} = \langle J\xi, \eta \rangle$$

Thus

$$\|u\|^2 \|v\|^2 - (\text{real}(\langle iu, v \rangle))^2 = \|\xi\|^2 \|\eta\|^2 - (\langle J\xi, \eta \rangle)^2 = \langle (\|\xi\|^2 1 - \llbracket J\xi, J\xi \rrbracket) \eta, \eta \rangle$$

Substituting in (3.83) we obtain (3.42).

(3) $\Leftrightarrow$ (4). Assume $\text{rank}(I(\xi)) = 2n - 1$ for all $\xi \neq 0$. A direct computation shows that $I(\xi)(J\xi) = 0$. Hence $J\xi$ is the only independent vector in $\ker(I(\xi))$. It follows there is $a = a(\xi) > 0$ so that

$$I(\xi) \geq a(\xi) P_{J\xi}^\perp$$

Note the $a(\xi)$ represent the smallest nonzero eigenvalue of $I(\xi)$ which must be the $2n - 1^{\text{th}}$. Using the homogeneity of degree 2 on the unit sphere of $H_{\mathbb{R}}$, we get $a(\xi) = \|\xi\|^2 a(\frac{\xi}{\|\xi\|}) \geq a_0 \|\xi\|^2$ which proves (3.42). Conversely, if (3.42) holds true, then $I(\xi)$ has rank at least $2n - 1$. Again since $J\xi$ is in the kernel of $I(\xi)$, it follows that $I(\xi)$ must be of rank exactly $2n - 1$. Q.E.D. $\square$

Proof of Corollary 3.5

(1) $\Leftrightarrow$ (2) follows from Theorem 3.1(2) and equation (3.51) and the fact that $\text{imag}(\langle u, v \rangle) = 0$ for all $u, v \in H'$.

(2) $\Leftrightarrow$ (5) follows from the relation

$$\sum_{k=1}^m |\langle \Phi'_k u, v \rangle|^2 = \sum_{k=1}^m \langle \llbracket \Phi'_k u, \Phi'_k u \rrbracket v, v \rangle$$

and

$$\sum_{k=1}^m \llbracket \Phi'_k u, \Phi'_k u \rrbracket = \sum_{k=1}^m \Phi'_k \llbracket u, u \rrbracket \Phi'_k$$

(4) $\Leftrightarrow$ (5) follows by the homogeneity of $I'(x)$ with respect to $\|x\|$.

(2) $\Leftrightarrow$ (3) follows by contradiction and homogeneity of $(u, v) \mapsto \sum_{k=1}^m |\langle \Phi'_k u, v \rangle|^2$ with respect to the product $\|u\| \|v\|$.

Q.E.D. □

4. PERFORMANCE BOUNDS ON RECONSTRUCTION ALGORITHMS

In this section we present two performance bounds applicable to any reconstruction algorithm. One bound is deterministic and is based on the constants a_0 introduced in Theorem 3.1. The second bound represents the Cramer-Rao Lower Bound for the stochastic model (2.3).

4.1. Lipschitz bounds of the inverse map. Consider the nonlinear map $\alpha : H \rightarrow \mathbb{R}^m$. We shall establish a deterministic performance bound for any inversion algorithm in terms of the Lipschitz bounds of the map:

$$(4.84) \quad \mathcal{A} : \mathcal{S}^{1,0} \rightarrow \mathbb{R}^m, \quad \mathcal{A}(\llbracket x, x \rrbracket) = \alpha(x)$$

Specifically we want to bound above and below the following expression:

$$(4.85) \quad R(x, y) = \frac{\|\mathcal{A}(\llbracket x, x \rrbracket) - \mathcal{A}(\llbracket y, y \rrbracket)\|^2}{\|\llbracket x, x \rrbracket - \llbracket y, y \rrbracket\|_1^2}$$

Since $\llbracket x, x \rrbracket - \llbracket y, y \rrbracket = \llbracket u, v \rrbracket \in \mathcal{S}^{1,1}$ for some $u, v \in H$ it follows:

$$(4.86) \quad \sup_{x, y \in H} R(x, y) = \sup_{u, v \in H} \frac{\sum_{k=1}^m |\langle F_k, \llbracket u, v \rrbracket \rangle|^2}{\|\llbracket u, v \rrbracket\|_1^2}$$

$$(4.87) \quad \inf_{x, y \in H} R(x, y) = \inf_{u, v \in H} \frac{\sum_{k=1}^m |\langle F_k, \llbracket u, v \rrbracket \rangle|^2}{\|\llbracket u, v \rrbracket\|_1^2}$$

These ratios can be further processed as follows

$$\frac{\sum_{k=1}^m |\langle F_k, \llbracket u, v \rrbracket \rangle|^2}{\|\llbracket u, v \rrbracket\|_1^2} = \frac{\langle I(\xi) \eta, \eta \rangle}{\|\xi\|^2 \langle P_{J\xi}^\perp \eta, \eta \rangle}$$

where $\xi = \mathbf{j}(u)$ and $\eta = \mathbf{j}(v)$. Since $I(\xi)\eta = I(\xi)P_{J\xi}^\perp\eta$ it follows:

$$\sup_{\xi, \eta \neq 0} \frac{\langle I(\xi)\eta, \eta \rangle}{\|\xi\|^2 \langle P_{J\xi}^\perp \eta, \eta \rangle} = \sup_{\xi \neq 0} \frac{\|I(\xi)\|}{\|\xi\|^2} = \max_{\xi \in H_{\mathbb{R}}, \|\xi\|=1} \|I(\xi)\|$$

and

$$\inf_{\xi, \eta \neq 0} \frac{\langle I(\xi)\eta, \eta \rangle}{\|\xi\|^2 \langle P_{J\xi}^\perp \eta, \eta \rangle} = a_0^{opt}$$

Thus we proved:

Theorem 4.1. *Assume the nonlinear map α is injective. Then the map $\mathcal{A} : \mathcal{S}^{1,0} \rightarrow \mathbb{R}^m$ defined in (4.84) is bi-Lipschitz between $(\mathcal{S}^{1,0}, \|\cdot\|_1)$ and $(\mathbb{R}^m, \|\cdot\|)$ with the Euclidian norm, and it has the upper Lipschitz bound*

$$(4.88) \quad B_0 = \max_{\xi \in H_{\mathbb{R}}, \|\xi\|=1} \sqrt{\|I(\xi)\|}$$

and the lower Lipschitz bound

$$(4.89) \quad A_0 = \sqrt{a_0^{opt}} = \sqrt{\min_{\xi \in H_{\mathbb{R}}, \|\xi\|=1} a_{n-1}(I(\xi))}$$

Specifically

$$(4.90) \quad A_0 \|\llbracket x, x \rrbracket - \llbracket y, y \rrbracket\|_1 \leq \|\mathcal{A}(\llbracket x, x \rrbracket) - \mathcal{A}(\llbracket y, y \rrbracket)\| \leq B_0 \|\llbracket x, x \rrbracket - \llbracket y, y \rrbracket\|_1$$

for all $u, v \in H$.

4.2. The Cramer-Rao Lower Bound (CRLB) . Consider now the noise model (2.3). We are interested in obtaining a lower bound for any unbiased estimator of x . The derivation of the CRLB in this paper coincides with the one presented in [11] (see Theorem 23 there). In turn this follows the recipe presented in [9]. We will just present the key steps of this derivation. Note that our derivation is canonical, that is basis independent.

Due to non-holomorphy of the nonlinear map α , the analysis is done in the realification space $H_{\mathbb{R}}$. We denote $\xi = \mathbf{j}(x)$ for the signal $x \in H$. The frame set if $\mathbb{F} = \{f_1, \dots, f_m\}$ and $F_k = \llbracket f_k, f_k \rrbracket \in \mathcal{S}^{1,0}(H)$ denote the measurement operators. Recall our notation $\Phi_k = \tau(F_k) = \llbracket \varphi_k, \varphi_k \rrbracket + \llbracket J\varphi_k, J\varphi_k \rrbracket \in \mathcal{S}^{2,0}(H_{\mathbb{R}})$. First we compute the Fisher information matrix associated to ξ . The likelihood for this problem is

$$(4.91) \quad p(y|\xi) = \frac{1}{(2\pi)^{m/2}\sigma^m} \exp\left(-\frac{1}{2\sigma^2} \|y - \alpha(x)\|^2\right) = \frac{1}{(2\pi)^{m/2}\sigma^m} \exp\left(-\frac{1}{2\sigma^2} \sum_{k=1}^m |y_k - \langle \Phi_k \xi, \xi \rangle|^2\right)$$

The Fisher information matrix is given by (see [43])

$$(4.92) \quad F(\xi) = \mathbb{E} [(\nabla_{\xi} \log(p(y|\xi)))(\nabla_{\xi} \log(p(y|\xi)))^T]$$

The canonical form of this operator is

$$(4.93) \quad F(\xi) = \mathbb{E} [\llbracket \nabla_{\xi} \log(p(y|\xi)), \nabla_{\xi} \log(p(y|\xi)) \rrbracket]$$

A little bit of algebra shows

$$(4.94) \quad F(\xi) = \frac{4}{\sigma^2} \sum_{k=1}^m \Phi_k \llbracket \xi, \xi \rrbracket \Phi_k = \frac{4}{\sigma^2} I(\xi)$$

In general the covariance of any unbiased estimator is bounded below by the inverse of the Fisher information matrix (operator). However in this case the Fisher information operator is not invertible. This fact simply expresses the statement that x is not identifiable from the measurements $y \in \mathbb{R}^m$ alone. As we know the nonlinear map α is not injective on H but instead it is injective on $\hat{H}$. The nonuniqueness on H is reflected in having a singular Fisher information matrix on $H_{\mathbb{R}}$. To solve this issue we need to fix the global phase factor. One solution is to fix a basis and decide that a particular component (say the last component) is real. Such an approach was taken in [11]. Here we propose a canonical solution to this normalization. An oracle provides us with a vector $z_0 \in H$, so that $\langle x, z_0 \rangle > 0$ is positive real. Assume z_0 is normalized $\|z_0\| = 1$. Note there are two pieces of information that can be extracted from here: First the fact that x is not orthogonal to z_0 ; in particular $x \neq 0$. Second, the global phase to recover x from its rank-1 operator $\llbracket x, x \rrbracket$ is uniquely determined by the fact that $\text{imag}(\langle x, z_0 \rangle) = 0$ and $\text{real}(\langle x, z_0 \rangle) > 0$.

Under this scenario we want to analyze the Fisher information operator obtained earlier. Let $\psi_0 = \mathbf{j}(z_0) \in H_{\mathbb{R}}$. We know

$$\langle \xi, \psi_0 \rangle = \text{real}(\langle x, z_0 \rangle) > 0, \quad \langle \xi, J\psi_0 \rangle = \text{imag}(\langle x, z_0 \rangle) = 0$$

Let Π denote the orthogonal projection onto the complement of $J\psi_0$,

$$(4.95) \quad \Pi : H_{\mathbb{R}} \rightarrow H_{\mathbb{R}}, \quad \Pi = 1 - \llbracket J\psi_0, J\psi_0 \rrbracket$$

and let $E \subset H_{\mathbb{R}}$ denote its range, $E = \{J\psi_0\}^{\perp}$. The oracle tells us $\xi \in E$. Hence the class of estimators for ξ should include only functions

$$(4.96) \quad \omega : \mathbb{R}^m \rightarrow \mathbf{j}^{-1}(E) = \{x \in H ; \langle x, z_0 \rangle > 0\}$$

In this case the appropriate Fisher information operator should be

$$(4.97) \quad \tilde{F}(\xi) = \Pi F(\xi) \Pi = \frac{4}{\sigma^2} \sum_{k=1}^m \Pi \Phi_k \llbracket \xi, \xi \rrbracket \Phi_k \Pi$$

The following lemma proves that under the scenario described here $\tilde{F}(\xi)$ is invertible on E .

Lemma 4.2. *Assume α is injective on $\hat{H}$ and $z_0 \in H$ is so that $\langle x, z_0 \rangle > 0$. Then*

$$(4.98) \quad \tilde{F}(\xi) = \Pi F(\xi) \Pi \geq \frac{4}{\sigma^2} a_0 |\langle x, z_0 \rangle|^2 \Pi$$

where a_0 is the same lower bound introduced in Theorem 3.1 whose optimal value is given by (3.44).

Proof

We know $I(\xi) \geq a_0 \|\xi\|^2 P_{J\xi}^\perp$ from Theorem 3.1 (4). We only need to show

$$\|\xi\|^2 \Pi P_{J\xi}^\perp \Pi \geq |\langle \xi, \psi_0 \rangle|^2 \Pi$$

and the inequality is tight. Without loss of generality we can assume $\|\xi\| = 1$ since all expressions are homogeneous in $\|\xi\|$. Then we need to show that for any $u \in E$, $\|u\| = 1$,

$$\langle P_{J\xi}^\perp u, u \rangle \geq |\langle \xi, \psi_0 \rangle|^2$$

This follows from

$$\inf_{\|u\|=1, u \in E} 1 - |\langle u, J\xi \rangle|^2 = 1 - \max_{\|u\|=1, u \in E} |\langle u, J\xi \rangle|^2 = 1 - \left| \left\langle \frac{\Pi J\xi}{\|\Pi J\xi\|}, J\xi \right\rangle \right|^2 = |\langle \xi, \psi_0 \rangle|^2$$

Note also the inequality is tight. Q.E.D. □

Thus we established that $\tilde{F}(\xi)$ is invertible on E . See also [11], Lemma 22, for a similar statement. Let H_{z_0} denote the following open set;

$$(4.99) \quad H_{z_0} = \{x \in H, \langle x, z_0 \rangle > 0\}$$

Recall an estimator $\omega : \mathbb{R}^m \rightarrow H$ is said *unbiased* if

$$(4.100) \quad \mathbb{E}[\omega(y)|x] = x$$

We can now state the main result of this section:

Theorem 4.3. *Assume the nonlinear map α is injective and fix a vector $z_0 \in H$. For any vector $x \in H_{z_0}$ the covariance operator of any unbiased estimator $\omega : \mathbb{R}^m \rightarrow H_{z_0}$ of x is bounded below by the Cramer-Rao Lower Bound (CRLB) given by*

$$(4.101) \quad \text{Cov}[\omega(y)] \geq \frac{\sigma^2}{4} \left(\sum_{k=1}^m \Phi_k \llbracket \xi, \xi \rrbracket \Phi_k \right)^\dagger$$

where $\dagger$ denotes the pseudoinverse operator. In particular the Mean Square Error of ω is bounded below by

$$(4.102) \quad \text{MSE}(\omega) = \mathbb{E}[\|x - \omega(y)\|^2 | x] \geq \frac{\sigma^2}{4} \text{tr} \left\{ \left(\sum_{k=1}^m \Phi_k \llbracket \xi, \xi \rrbracket \Phi_k \right)^\dagger \right\}$$

The previous Lemma allows us to predict an upper bound for the MSE of any efficient estimator (that is an unbiased estimator that achieves CRLB):

Corollary 4.4. *Assume $\omega : \mathbb{R}^m \rightarrow H_{z_0}$ is an unbiased estimator that achieves the CRLB (4.101). Then its Mean-Square Error is bounded above by*

$$(4.103) \quad \text{MSE}(\omega) = \mathbb{E}[\|x - \omega(y)\|^2 | x] \leq \frac{(2n-1)\sigma^2}{4a_0^{\text{opt}} |\langle x, z_0 \rangle|^2}$$

5. THE REGULARIZED ITERATIVE LEAST-SQUARE ALGORITHM

Consider the additive noise model in (2.3). Our data is the vector $y \in \mathbb{R}^m$. Our goal is to find a $x \in H$ that minimizes $\|y - \alpha(x)\|$, where we use the Euclidian norm. Set

$$(5.104) \quad J_0(X) = \sum_{k=1}^m |y_k - \langle X f_k, f_k \rangle|^2, \quad J_0 : \text{Sym}(H) \rightarrow \mathbb{R}$$

and notice $J_0(\llbracket x, x \rrbracket) = \|y - \alpha(x)\|^2$. The least-square error minimizer represents the Maximum Likelihood Estimator (MLE) when the noise is Gaussian. In this section we discuss an optimization algorithm for this criterion.

Consider now $J_0 = \|y - \mathcal{A}(X)\|^2$ restricted to $\mathcal{S}^{1,0}$, which is an analytic manifold. Consider a current point $X^t = \llbracket x^t, x^t \rrbracket$ in an iterative process. Then a descent direction can be thought of as a vector in the tangent space to the manifold. According to Lemma 3.12 (1), the tangent space at X^t is given by operators of the form $\llbracket x^t, \delta \rrbracket$. Since $X^t + \llbracket x^t, \delta \rrbracket = \llbracket x^t, x^t + \delta \rrbracket \in \mathcal{S}^{1,1}$, one would need to project $X^t + \llbracket x^t, \delta \rrbracket$ back into $\mathcal{S}^{1,0}$ and choose direction δ that minimizes (or at least decreases) $J_0(P(X^t + \llbracket x^t, \delta \rrbracket))$, where P is the (nonlinear) projection in $\text{Sym}(H)$ onto $\mathcal{S}^{1,0}$. However since J is well defined on $\mathcal{S}^{1,1}$ we choose to optimize δ without projecting back onto $\mathcal{S}^{1,0}$. Thus we obtain the iterative process:

$$x^{t+1} = \operatorname{argmin}_u J_0(\llbracket x^t, u \rrbracket)$$

However this process is not robust to noise, the main reason being ill-conditioning and multiple local minima of J_0 on $\mathcal{S}^{1,0}$. Instead we choose to regularize this process and thus to introduce a different optimization criterion.

Consider the following functional

$$(5.105) \quad J : H \times H \times \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$$

$$J(u, v, \lambda, \mu) = \sum_{k=1}^m |y_k - \frac{1}{2}(\langle u, f_k \rangle \langle f_k, v \rangle + \langle v, f_k \rangle \langle f_k, u \rangle)|^2 + \lambda \|u\|^2 + \mu \|u - v\|^2 + \lambda \|v\|^2.$$

Our ultimate goal is to minimize $J_0(u) = \|y - \alpha(u)\|^2 = J(u, u, 0, \mu)$ over u , for some (and hence any) value of $\mu \in \mathbb{R}^+$. Our strategy is based on the following iterative process:

Step 0. we initialize x^0 as the global optimal solution for a specific pair (λ_0, μ_0) .

Step 1. Repeat:

$$(5.106) \quad x^{t+1} = \operatorname{argmin}_u J(u, x^t, \lambda_t, \mu_t)$$

Update λ_{t+1}, μ_{t+1} according to a specific policy;

Step 3. Stop when some tolerance level is achieved.

As we describe below the update (5.106) can be tweaked to achieve a more robust behavior (see (5.124)).

5.1. Initialization. Consider the regularized least-square problem:

$$\min_u J(u, u, \lambda, 0) = \min_u \|y - \alpha(u)\|^2 + 2\lambda \|u\|^2$$

Note the following relation

$$\begin{aligned}
 J(u, u, \lambda, 0) &= \|y\|^2 + 2\lambda\|u\|^2 - 2 \sum_{k=1}^m y_k |\langle u, f_k \rangle|^2 + \sum_{k=1}^m |\langle u, f_k \rangle|^4 \\
 (5.107) \quad &= \|y\|^2 + 2\langle (\lambda I - Q)u, u \rangle + \sum_{k=1}^m |\langle u, f_k \rangle|^4
 \end{aligned}$$

where

$$(5.108) \quad Q = \sum_{k=1}^m y_k \llbracket f_k, f_k \rrbracket = \sum_{k=1}^m y_k F_k$$

For $\lambda > \|Q\|$ the optimal solution is $u = 0$. Note that if $Q \leq 0$ (as a quadratic form) then the optimal solution of $\min_u \|y - \varphi(u)\|^2$ is $u = 0$. Consequently we assume the largest eigenvalue of Q is positive. As λ decreases the optimizer remains small. Hence we can neglect the forth order term in u in the expansion above and obtain:

$$J(u, u, \lambda, 0) \approx \|y\|^2 + 2\langle (\lambda I - Q)u, u \rangle$$

Thus the critical value of λ for which we may get a nonzero solution is $\lambda = \max \text{eig}(Q)$ is the maximum eigenvalue of Q . Let us denote by a_1 this (positive) eigenvalue and e_1 its associated normalized eigenvector. This suggests to initialize $\lambda = \rho a_1$ for some $0 < \rho \leq 1$ and $x^0 = \beta e_1$, for some scalar β . Substituting into (5.107) we obtain

$$J(\beta e_1, \beta e_1, \rho a_1, 0) = \|y\|^2 - 2(1 - \rho)a_1\beta^2 + \left(\sum_{k=1}^m |\langle e_1, f_k \rangle|^4\right)\beta^4$$

For fixed ρ , the minimum over β is achieved at

$$(5.109) \quad \beta_0 = \sqrt{\frac{(1 - \rho)a_1}{\sum_{k=1}^m |\langle e_1, f_k \rangle|^4}}, \quad x^0 = \beta e_1$$

The parameter μ controls the step size at each iteration. The larger the value the smaller the step. On the other hand, a small value of this parameter may produce an unstable behavior of the iterates. In our implementation we use the same initial value for both λ and μ :

$$(5.110) \quad \mu_0 = \lambda_0 = \rho a_1$$

5.2. Iterations. Minimization (5.106) is performed in the space $H_{\mathbb{R}}$. Let $\xi^t = \mathbf{j}(x^t)$ and $\eta = \mathbf{j}(u)$. Then

$$(5.111) \quad J(u, x^t, \lambda_t, \mu_t) = \sum_{k=1}^m |\langle \llbracket \Phi_k \xi^t, \Phi_k \xi^t \rrbracket \eta, \eta \rangle - y_k|^2 + \lambda_t \|\eta\|^2 + \mu_t \|\eta - \xi^t\|^2 + \lambda_t \|\xi^t\|^2$$

Note the criterion is quadratic in η . The unique minimum is given by solving the linear equation:

$$(5.112) \quad \left(\sum_{k=1}^m \Phi_k \llbracket \xi^t, \xi^t \rrbracket \Phi_k + (\lambda_t + \mu_t) 1 \right) \xi^{t+1} = \left(\sum_{k=1}^m y_k \Phi_k + \mu_t 1 \right) \xi^t$$

for ξ^{t+1} . In our implementations we decrease (λ_t, μ_t) but we limit μ_t to a minimum value. Thus our adaptation policy is

$$(5.113) \quad \lambda_{t+1} = \gamma \lambda_t$$

$$(5.114) \quad \mu_{t+1} = \max(\gamma \mu_t, \mu^{min})$$

where $0 < \gamma < 1$ is the rate parameter.

5.3. Stopping Criterion. In our implementation we repeat the iterations until λ reaches a preset value λ^{min} . As proved later in this section, the error is linearly dependent on λ .

Alternatively, one can stop the iterations once the modeling error becomes of comparable to noise variance. Specifically, a stopping criterion would be

$$(5.115) \quad \sum_{k=1}^m |y_k - |\langle x^t, f_k \rangle|^2|^2 \leq \kappa m \sigma^2$$

where $\kappa \gtrsim 1$, for instance $\kappa = 3$.

5.4. Convergence and Optimality. Consider the following three functionals $J_1, J_2, J_3 : \text{Sym}(H) \times \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}$ defined by

$$(5.116) \quad J_1(X, \lambda, \mu) = \sum_{k=1}^m |y_k - \langle X, F_k \rangle|^2 + 2(\lambda + \mu) \|X\|_1 - 2\mu \text{tr}\{X\}$$

$$(5.117) \quad J_2(X, \lambda, \mu) = \sum_{k=1}^m |y_k - \langle X, F_k \rangle|^2 + 2\lambda a_{max}(X) - 4\mu a_{min}(X)$$

$$(5.118) \quad J_3(X, \lambda, \mu) = \sum_{k=1}^m |y_k - \langle X, F_k \rangle|^2 + 2\lambda \|X\|_1 - 2(2\lambda - \mu) a_{min}(X)$$

where $a_{max}(X)$ and $a_{min}(X)$ are the maximum and minimum eigenvalue of X , respectively. We can prove the following result:

Lemma 5.1. (1) *When restricted to $\mathcal{S}^{1,1}$ the three criteria coincide:*

$$(5.119) \quad J_1(X, \lambda, \mu) = J_2(X, \lambda, \mu) = J_3(X, \lambda, \mu), \quad \forall X \in \mathcal{S}^{1,1}, \lambda, \mu \in \mathbb{R}.$$

(2) *On $\text{Sym}(H)$, the three criteria J_1, J_2, J_3 are convex.*

(3) *The minimum value of J_1, J_2, J_3 on $\mathcal{S}^{1,1}$ coincides with the minimum value of J on $H \times H$:*

$$(5.120) \quad \min_{X \in \mathcal{S}^{1,1}} J_1(X, \lambda, \mu) = \min_{X \in \mathcal{S}^{1,1}} J_2(X, \lambda, \mu) = \min_{X \in \mathcal{S}^{1,1}} J_3(X, \lambda, \mu) = \min_{u, v \in H} J(u, v, \lambda, \mu)$$

for any $\lambda, \mu \geq 0$. Any minimizer $\hat{X} \in \mathcal{S}^{1,1}$ for J_1, J_2, J_3 and $(\hat{u}, \hat{v})$ for J satisfy

$$(5.121) \quad \hat{X} = \llbracket \hat{u}, \hat{v} \rrbracket, \quad \|\hat{u}\| = \|\hat{v}\|, \quad \text{imag}(\langle u, v \rangle) = 0$$

(4) Restricted to $\mathcal{S}^{1,0}$ all four criteria coincide:

$$(5.122) \quad J(u, u, \lambda, \mu) = J_1(\llbracket u, u \rrbracket, \lambda, \mu) = J_2(\llbracket u, u \rrbracket, \lambda, \mu) = J_3(\llbracket u, u \rrbracket, \lambda, \mu) = \|y - \alpha(u)\|^2 + 2\lambda\|u\|^2$$

and are independent of μ .

Proof

(1) and (2) can be checked directly.

(3) and (4) follow from the following relation:

$$(5.123) \quad \begin{aligned} J(u, v, \lambda, \mu) - J_1(\llbracket u, v \rrbracket, \lambda, \mu) &= (\lambda + \mu)[(\|u\| - \|v\|)^2 + 2\|u\|\|v\| - 2\sqrt{\|u\|^2\|v\|^2 - (\text{imag}(\langle u, v \rangle))^2}] \\ &\geq (\lambda + \mu)(\|u\| - \|v\|)^2 \end{aligned}$$

that follows from (3.67) and (3.71).

Remark 5.2. The criterion J_2 shows that the two regularization terms $\lambda(\|u\|^2 + \|v\|^2)$ and $\mu\|u - v\|^2$ have different effects on the optimizer: the larger the parameter μ the closer the lower eigenvalue is to zero; hence the closer the optimizer $\hat{X}$ is to a rank-1 operator; on the other hand, the larger the parameter λ the larger the cost of the $\mathcal{S}^{1,0}$ component in $\hat{X}$; hence $\|\hat{X}\|$ remains bounded.

Remark 5.3. In the optimization problem $\inf_{u,v \in H} J(u, v, \lambda, \mu)$ the constraint is convex but the criterion is not jointly convex in (u, v) . It is however convex in each individual variable u and v . On the other hand in the optimization problems $\inf_{X \in \mathcal{S}^{1,1}} J_s(X, \lambda, \mu)$, $1 \leq s \leq 3$, the criterion is convex in X , but the underlying constraint $X \in \mathcal{S}^{1,1}$ does not define a convex set.

The optimization procedure outlined in the previous subsection describes the following mechanism. Consider a path $(x^t)_{t \geq 0}$ in H . Set

$$X^{t+1} = \llbracket x^t, x^{t+1} \rrbracket = \frac{1}{2} (\langle \cdot, x^{t+1} \rangle x^t + \langle \cdot, x^t \rangle x^{t+1})$$

Thus a trajectory $(x^t)_{t \geq 0}$ in H is mapped into a trajectory $(X^t)_{t \geq 0}$ in $\mathcal{S}^{1,1}$. Then the algorithm chooses X^{t+1} along a tangent direction to $\mathcal{S}^{1,1}$ at $\llbracket x^t, x^{t-1} \rrbracket$, namely a direction of the form $\llbracket x^t, u \rrbracket$ for some $u \in H$ that is of maximum descent for J_s , $1 \leq s \leq 3$. Since the algorithm performance is completely characterized by the sequence $(X^t)_{t \geq 0}$, and since for a given $X \in \mathcal{S}^{1,1}$ the minimum of $J(u, v, \lambda, \mu)$ over $u, v \in H$ subject to $\llbracket u, v \rrbracket = X$ is achieved at a pair (u, v) so that $\|u\| = \|v\|$ and $\text{imag}(\langle u, v \rangle) = 0$, we chose to rescale the vector x^{t+1} to achieve norm $\sqrt{\|\xi^t\| \|\xi^{t+1}\|}$. Thus:

$$(5.124) \quad x^{t+1} = \sqrt{\frac{\|\xi^t\|}{\|\xi^{t+1}\|}} \mathbf{j}^{-1}(\xi^{t+1})$$

5.5. Robustness to noise and the effect of regularization. We this subsection we make a stronger assumption of injectivity:

Assumption A For every $x \in H$ there is a unique $X \in \mathcal{S}^{1,0}$ so that $\mathcal{A}(X) = \mathcal{A}(\llbracket x, x \rrbracket)$, namely $X = \llbracket x, x \rrbracket$.

A heuristic argument (that can be made more precise, but we do not intend to do so here) suggests that generically this assumption is satisfied for frames of redundancy 6 or more (that is for $m \geq 6n$).

This assumption turns out to be equivalent to a stability bound that we describe next

Lemma 5.4. *The following are equivalent:*

- (1) *The frame $\mathbb{F}$ satisfies hypothesis A.*
- (2) $\ker \mathcal{A} \cap \mathcal{S}^{2,1} = \{0\}$.
- (3) *There is a positive constant $A_3 > 0$ so that*

$$(5.125) \quad A_3 \|X - Y\|_1^2 \leq \|\mathcal{A}(X) - \mathcal{A}(Y)\|^2$$

for all $X \in \mathcal{S}^{1,1}$ and $Y \in \mathcal{S}^{1,0}$.

Proof

(1) $\Leftrightarrow$ (2) is immediate once we notice that $\mathcal{S}^{1,0} - \mathcal{S}^{1,1} = \mathcal{S}^{2,1}$ cf. Lemma 3.6 (5).

(3) $\Rightarrow$ (2) is immediate.

(2) $\Rightarrow$ (3). Since $S = \{W \in \mathcal{S}^{2,1} \mid \|W\|_1 = 1\}$ is compact, it follows

$$(5.126) \quad A_3 = \inf_{W \in S} \|\mathcal{A}(W)\|^2 = \|\mathcal{A}(W_0)\|^2 > 0$$

for some $W_0 \in \mathcal{S}^{2,1}$, $W_0 \neq 0$. Then since any $Z \in \mathcal{S}^{2,1}$ can be written as $Z = \|Z\|_1 W$, with $W = \frac{1}{\|Z\|_1} Z \in S$,

$$\|\mathcal{A}(Z)\|^2 = \|Z\|_1^2 \|\mathcal{A}(W)\|^2 \geq A_3 \|Z\|_1^2$$

Then (5.125) follows from noticing that $Y - X \in \mathcal{S}^{2,1}$. Q.E.D.

Now we can state the main stability result of the estimators described in this section.

Theorem 5.5. *Fix $\mu \geq 2\lambda$, $x \in H_{z_0}$ and let $y = \alpha(x) + \nu$. Assume the optimization procedure finds a pair $u, v \in H$ so that $J(u, v, \lambda, \mu) \leq J(x, x, \lambda, \mu)$. Then*

$$(5.127) \quad \|\llbracket u, v \rrbracket - \llbracket x, x \rrbracket\|_1 \leq \frac{2\lambda}{A_3} + 2\sqrt{\frac{\lambda^2}{A_3^2} + \|\nu\|^2} \leq 6\frac{\lambda}{A_3} + 4\|\nu\|$$

Let $\llbracket u, v \rrbracket = a_1 \llbracket e_1, e_1 \rrbracket - a_2 \llbracket e_2, e_2 \rrbracket$, with $a_1, a_2 \geq 0$ and $\{e_1, e_2\}$ orthonormal set, be its spectral decomposition. Assume an oracle provides the global phase $e^{i\varphi_0} = \frac{\langle x, e_1 \rangle}{|\langle x, e_1 \rangle|}$. Set

$$(5.128) \quad \hat{x} = e^{i\varphi_0} \sqrt{a_1} e_1$$

Then

$$(5.129) \quad \|x - \hat{x}\|^2 \leq \|\llbracket u, v \rrbracket - \llbracket x, x \rrbracket\|_1 + |a_2(\llbracket u, v \rrbracket)|$$

If, additionally, $J(u, v, \lambda, \mu) \leq J(0, 0, \lambda, \mu)$ then

$$(5.130) \quad \|x - \hat{x}\|^2 \leq 6\frac{\lambda}{A_3} + 4\|\nu\| + \frac{\|\nu\|^2}{4\mu}$$

Proof

The proof follows from the Lipschitz bound (5.125). Let $X = \llbracket u, v \rrbracket$ and $Y = \llbracket x, x \rrbracket$. Note that

$$J_3(X, \lambda, \mu) \leq J(u, v, \lambda, \mu) \leq J(x, x, \lambda, \mu) = J_3(\llbracket x, x \rrbracket, \lambda, \mu)$$

Explicitely this means

$$\|\mathcal{A}(X - Y) - \nu\|^2 + 2\lambda\|X\|_1 - 2(2\mu - \lambda)a_{\min}(X) \leq \|\nu\|^2 + 2\lambda\|Y\|_1$$

Then

$$A_3\|X - Y\|_1^2 \leq \|\mathcal{A}(X - Y)\|^2 \leq (\|\mathcal{A}(X - Y) - \nu\| + \|\nu\|)^2 \leq 2\|\mathcal{A}(X - Y) - \nu\|^2 + 2\|\nu\|^2$$

Substituting into the previous inequality we obtain

$$\|X - Y\|_1^2 - \frac{4\lambda}{A_3}\|X - Y\|_1 - 4(\|\nu\|^2 - (2\lambda - \mu)a_{\min}(X)) \leq 0$$

Solving for $\|X - Y\|_1$ we obtain

$$\|X - Y\|_1 \leq \frac{2\lambda}{A_3} + 2\sqrt{\frac{\lambda^2}{A_3^2} + \|\nu\|^2 - (2\lambda - \mu)a_{\min}(X)}$$

which together with $\mu \geq 2\lambda$ and $a_{\min}(X) \leq 0$ for any $X \in \mathcal{S}^{1,1}$ proves the first inequality in (5.127). The second inequality in (refeq:robust) follows from $\sqrt{a^2 + b^2} \leq a + b$ for any $a, b \geq 0$.

The second part of the Theorem is obtained as follows. First note $\llbracket \hat{x}, \hat{x} \rrbracket = a_1\llbracket e_1, e_1 \rrbracket$. Hence

$$\|\llbracket \hat{x}, \hat{x} \rrbracket - \llbracket x, x \rrbracket\|_1 \leq \|\llbracket u, v \rrbracket - \llbracket x, x \rrbracket\|_1 + |a_2(\llbracket u, v \rrbracket)|$$

Next we show that $\|x - \hat{x}\|^2 \leq \|\llbracket \hat{x}, \hat{x} \rrbracket - \llbracket x, x \rrbracket\|_1$, from where (5.129) follows. Let $T = \llbracket \hat{x}, \hat{x} \rrbracket - \llbracket x, x \rrbracket \in \mathcal{S}^{1,1}$. The two (possibly) nonzero eigenvalues are obtained from solving the system (3.74). Explicitely we obtain:

$$(5.131) \quad a_{\pm} = \frac{1}{2}\text{tr}\{T\} \pm \frac{1}{2}\sqrt{2\text{tr}\{T^2\} - (\text{tr}\{T\})^2}$$

Thus

$$\|T\|_1 = a_+ - a_- = \sqrt{2\text{tr}\{T^2\} - (\text{tr}\{T\})^2}$$

Direct computations show:

$$\text{tr}\{T\} = \|\hat{x}\|^2 - \|x\|^2, \quad \text{tr}\{T^2\} = \|\hat{x}\|^4 + \|x\|^4 - 2|\langle \hat{x}, x \rangle|^2$$

Thus:

$$\|T\|_1 = (\|\hat{x}\|^2 + \|x\|^2)^2 - 4|\langle x, \hat{x} \rangle|^2$$

Recall $\hat{x}$ has the global phase so that $\langle \hat{x}, x \rangle \geq 0$ is real and nonnegative. Thus $|\langle \hat{x}, x \rangle|^2 = (\langle \hat{x}, x \rangle)^2$ and

$$\|x - \hat{x}\|^4 = \|T\|_1^2 - 4\langle \hat{x}, x \rangle \cdot \|x - \hat{x}\|^2 \leq \|T\|_1^2$$

which shows $\|x - \hat{x}\|^2 \leq \|\llbracket \hat{x}, \hat{x} \rrbracket - \llbracket x, x \rrbracket\|_1$ and thus (5.129).

Finally, (5.130) follows from using the last inequality (5.127) together with a bound for $|a_2|$ from

$$4\mu|a_2| \leq J_3(X, \lambda, \mu) \leq J_3(0, \lambda, \mu) = \|\nu\|^2.$$

Q.E.D. □

Remark 5.6. *The assumption $J(u, v, \lambda, \mu) < J(0, 0, \lambda, \mu)$ is not critical. A different upper bound than (5.130) can be obtained from $-4\mu a_2(X) \leq J_3(Y, \lambda, \mu) = \|\nu\|^2 + 2\lambda\|x\|^2$:*

$$(5.132) \quad \|x - \hat{x}\|^2 \leq 6\frac{\lambda}{A_3} + 4\|\nu\| + \frac{\|\nu\|^2}{4\mu} + \frac{\lambda\|x\|^2}{2\mu}$$

which can be made small (to the order of noise level) by choosing a large μ and small λ .

Remark 5.7. *Note the results does not require to find the global minimum, but only an estimate that brings the criterion J below the level achieved by the true signal. On the other hand the result is not unexpected. Similar bounds are known for MLE estimates.*

6. NUMERICAL ANALYSIS

In this section we present numerical simulations for the Regularized Iterative Least-Square algorithm presented in the previous section.

We generated random frames or redundancy $\frac{m}{n} = 4, 6$, and 8, as well as complex random signals x of size $n = 100$. All these vectors (frame and signal) are drawn from $\mathbb{N}(0, I)$. Then we scale the frame vectors to have norm 1. We set the first component of x to be a positive real, and so the global phase becomes uniquely determined. The algorithm stopped when λ_t reached a preset value. In these simulations we choose $\lambda^{\min} = 0.01$, $\mu^{\min} = 1$ and rate $\gamma = 0.8$ in (5.113, 5.114).

The magnitude square of signal coefficients $\alpha(x)$ is perturbed additively by Gaussian noise with variance σ^2 to achieve a fixed signal-to-noise-ratio defined as

$$SNR = \frac{\sum_{k=1}^m |\langle x, f_k \rangle|^4}{m\sigma^2}, \quad SNR_{dB} = 10 \log_{10}(SNR) \text{ [dB]}$$

We vary SNRdB over 15 values in 5dB increments from -30dB to +40dB. We average algorithm performance over 100 noise realizations.

Figure 1 includes the mean-square error averaged over 100 noise realizations for one fixed realization of x , and the C-R lower bound.

In Figure 2 we plot the bias and variance components of the mean-square error for the same results in Figure 1. Note the bias is relatively small. The bulk of mean-square error is due to estimation variance.

The mean number of iterations varied between 35 and 50, with a higher value for lower SNR (-30dB) and a smaller value for higher SNR (40dB).

For $m = 800$ and three values of SNR (-30dB, 0dB, and 40dB) Figures 3-5 plot traces of a particular noise realization (different for each SNR). In each figure, we plot four characteristics: top graph plots the estimation error of the clean signal $10 \log_{10} \|x^t - x\|^2$; second plot presents the smallest eigenvalue of $X^t = \llbracket x^t, x^{t-1} \rrbracket$ (which is negative); third plot contains the current estimation error of the clean rank-one, that is $10 \log_{10} \|X^t - \llbracket x, x \rrbracket\|_2^2$ (Frobenius norm); bottom graph plots $J_0 = 10 \log_{10} \|y - \mathcal{A}(X^t)\|^2$.

We also analyzed the algorithm sensitivity to initialization. If instead of using the eigen-pair (5.109, 5.110) we initialize by a random vector x^0 together with the eigenvalue (5.110). Results for the three values of redundancy are presented in Figure 6 for 0dB to 40dB range of SNR .

7. CONCLUSIONS

Novel necessary conditions for complex valued signal reconstruction from magnitudes of frame coefficients have been presented. Deterministic stability bounds (Lipschitz constants) and stochastic performance bounds (Cramer-Rao lower bound) have been presented. The entire analysis has been done canonically, that is independent of a particular choice of basis. Then an optimization algorithm based on the least-square error has been proposed and analyzed. The algorithm performance has been compared to the theoretical lower bound given by the Cramer-Rao inequality. Remarkably the algorithm performs very well on a large range of SNR. In particular, for high SNR, it seems to converge to the correct signal every time. This behavior suggests the algorithm presented here is able to track the global minimum of (5.105) very well. A future study shall analyze this tracking hypothesis.

ACKNOWLEDGMENTS

The author was partially supported by NSF under DMS-1109498 grant. The author is also grateful to the Erwin Schrödinger Institute for the hospitality shown during the special workshop on "Phase Retrieval" in October 2012. Some of the results obtained here were presented at that workshop and later at the Workshop on "Phaseless Reconstruction", UMD, February 2013. The author also thanks Bernhard Bodmann, Jameson Cahill, Martin Ehler, Boaz Nadler, Oren Raz, and Yang Wang for fruitful discussions.

REFERENCES

- [1] B. Alexeev, A. S. Bandeira, M. Fickus, D. G. Mixon, *Phase retrieval with polarization*, arXiv submission, arXiv:1210.7752
- [2] B. Alexeev, J. Cahill, D. G. Mixon, *Full spark frames*, J. Fourier Anal. Appl. 18 (2012) 11671194.
- [3] D. M. Appleby, *Symmetric informationally complete-positive operator valued measures and the extended Clifford group*, J. Math. Phys. **46** (2005), no. 5, 052107, 29.
- [4] R. Balan, P. Casazza, D. Edidin, *On signal reconstruction without phase*, Appl.Comput.Harmon.Anal. **20** (2006), 345–356.
- [5] R. Balan, P. Casazza, D. Edidin, *Equivalence of Reconstruction from the Absolute Value of the Frame Coefficients to a Sparse Representation Problem*, IEEE Signal.Proc.Letters, **14** (5) (2007), 341–343.

- [6] R. Balan, B. Bodmann, P. Casazza, D. Edidin, *Painless reconstruction from Magnitudes of Frame Coefficients*, J. Fourier Anal. Appl., **15** (4) (2009), 488–501.
- [7] R. Balan, *A Nonlinear Reconstruction Algorithm from Absolute Value of Frame Coefficients for Low Redundancy Frames*, Proceedings of SampTA Conference, Marseille, France May 2009.
- [8] R. Balan, *On Signal Reconstruction from Its Spectrogram*, Proceedings of the CISS Conference, Princeton NJ, May 2010.
- [9] R. Balan, *Reconstruction of Signals from Magnitudes of Frame Representations*, arXiv submission , arXiv:1207.1134
- [10] R. Balan, Y. Wang, , in preparation 2013.
- [11] A. S. Bandeira, J. Cahill, D. Mixon, A. A. Nelson, *Saving phase: Injectivity and Stability for phase retrieval*, arXiv submission , arXiv: 1302.4618
- [12] S. Bandyopadhyay, P. O. Boykin, V. Roychowdhury, and F. Vatan, *A new proof for the existence of mutually unbiased bases*, Algorithmica **34** (2002), no. 4, 512–528, Quantum computation and quantum cryptography.
- [13] B. G. Bodmann, private communications, October 2012 and March 2013.
- [14] B. G. Bodmann, N. Hammen, *Stable Phase Retrieval with Low-Redundancy Frames*, arXiv submission:1302.5487v1
- [15] B. G. Bodmann, *Optimal linear transmission by loss-insensitive packet encoding*, Appl. Comput. Harmon. Anal. **22** (2007), 274–285.
- [16] B. G. Bodmann and V. I. Paulsen, *Frames, graphs and erasures*, Linear Algebra Appl. **404** (2005), 118–146.
- [17] B. G. Bodmann, , arXiv submission , arXiv:1302.
- [18] P. O. Boykin, M. Sitharam, P. H. Tiep, and P. Wocjan *Mutually unbiased bases and orthogonal decompositions of Lie algebras*, E-print: arXiv.org/quant-ph/0506089, to appear in Quantum Information and Computation, 2007.
- [19] A. R. Calderbank, P. J. Cameron, W. M. Kantor, and J. J. Seidel, *Z_4 -Kerdock codes, orthogonal spreads, and extremal Euclidean line-sets*, Proc. London Math. Soc. (3) **75** (1997), no. 2, 436–480.
- [20] E. Candès, T. Strohmer, V. Voroninski, *PhaseLift: Exact and Stable Signal Recovery from Magnitude Measurements via Convex Programming*, to appear in Communications in Pure and Applied Mathematics (2011)
- [21] E. Candès, Y. Eldar, T. Strohmer, V. Voloninski, *Phase Retrieval via Matrix Completion Problem*, arXiv submission, arXiv:1109.0573
- [22] E. J. Candès, X. Li, *Solving quadratic equations via PhaseLift when there are about as many equations as unknowns*, arXiv submission , arXiv:1208.6247
- [23] P. Casazza, *The art of frame theory*, Taiwanese J. Math., (2) **4** (2000), 129–202.
- [24] P. G. Casazza and M. Fickus, *Fourier transforms of finite chirps*, EURASIP J. Appl. Signal Process. (2006), no. Frames and overcomplete representations in signal processing, communications, and information theory, Art. ID 70204, 1–7.
- [25] P. Casazza and J. Kovačević, *Equal-norm tight frames with erasures*, Adv. Comp. Math. **18** (2003), 387–430.
- [26] P. G. Casazza and G. Kutyniok, *Frames of subspaces*, Wavelets, frames and operator theory, Contemp. Math., vol. 345, Amer. Math. Soc., Providence, RI, 2004, pp. 87–113.
- [27] P. J. Cameron and J. J. Seidel, *Quadratic forms over $GF(2)$* , Indag. Math. **35** (1973), 1–8.
- [28] P. Delsarte, J. M. Goethals, and J. J. Seidel, *Spherical codes and designs*, Geometriae Dedicata **6** (1977), no. 3, 363–388.
- [29] L. Demanet, P. Hand, *Stable optimizationless recovery from phaseless linear measurements*, arXiv submission , arXiv:1208.1803v1

- [30] Y. Eldar, S. Mendelson, *Phase Retrieval: Stability and Recovery Guarantees*, arXiv submission, arXiv:1211.0872v1
- [31] A. Fannjiang, *Absolute uniqueness in phase retrieval with random illumination*, Inverse Probl. **28** (2012).
- [32] J. Finkelstein, *Pure-state informationally complete and “really” complete measurements*, Phys. Rev. A **70** (2004), no. 5, doi:10.1103/PhysRevA.70.052107
- [33] C. Godsil and A. Roy, *Equiangular lines, mutually unbiased bases and spin models*, 2005, E-print: arxiv.org/quant-ph/0511004.
- [34] V. K. Goyal, J. Kovačević, and J. A. Kelner, *Quantized frame expansions with erasures*, Appl. Comp. Harm. Anal. **10** (2001), 203–233.
- [35] P. R. Halmos, **Finite-Dimensional Hilbert Space**, Springer-Verlag, 1974.
- [36] P. Hand, *Conditions for Existence of Dual Certificates in Rank One Semidefinite Problems*, arXiv submission, arXiv:1303.1598v2
- [37] M. H. Hayes, J. S. Lim, and A. V. Oppenheim, *Signal Reconstruction from Phase and Magnitude*, IEEE Trans. ASSP **28**, no.6 (1980), 672–680.
- [38] T. Heinosaari, L. Mazzarella, M. M. Wolf, *Quantum Tomography under Prior Information*, arXiv:1109.5478v1 [quant-ph], 26 Sept 2011.
- [39] S. G. Hoggar, *t*-designs in projective spaces, Europ. J. Comb. **3** (1982), 233–254.
- [40] S. D. Howard, A. R. Calderbank, and W. Moran, *The finite Heisenberg-Weyl groups in radar and communications*, EURASIP J. Appl. Signal Process. (2006), no. Frames and overcomplete representations in signal processing, communications, and information theory, Art. ID 85685, 1–12.
- [41] R. Holmes and V. I. Paulsen, *Optimal frames for erasures*, Lin. Alg. Appl. **377** (2004), 31–51.
- [42] P. Jaming, *Uniqueness results for the phase retrieval problem of fractional Fourier transforms of variable order*, arXiv submission, arXiv:1009.3418
- [43] S. M. Kay, **Fundamentals of Statistical Signal Processing. I. Estimation Theory**, Prentice Hall PTR, 18th Printing, 2010.
- [44] J. Kovačević, P. L. Dragotti, and V. K. Goyal, *Filter bank frame expansions with erasures*, IEEE Trans. Inform. Theory **48** (2002), 1439–1450.
- [45] A. Klappenecker and M. Rötteler, *Mutually unbiased bases are complex projective 2-designs*, Proc. Int. Symp. on Inf. Theory, IEEE, 2005, pp. 1740–1744.
- [46] T.-Y. Lam, **Algebraic Theory of Quadratic Forms**, W A Benjamin, 1973.
- [47] R. G. Lane, W. R. Freight, and R. H. T. Bates, *Direct Phase Retrieval*, IEEE Trans. ASSP **35**, no.4 (1987), 520–526.
- [48] P. W. H. Lemmens and J. J. Seidel, *Equiangular lines*, J. Algebra **24** (1973), 494–512.
- [49] Yu. I. Lyubich, *On tight projective designs*, E-print: www.arXiv.org/math/0703526, 2007.
- [50] K. H. Mayer, *Elliptische Differentialoperatoren und Ganzzahligkeitssatze für charakteristische Zahlen*, Topology **4** (1965) 295–313.
- [51] R. J. Milgram, *Immersing projective spaces*, Ann. Math. **85** (1967) 473–482.
- [52] A. Mukherjee, *Embedding complex projective spaces in Euclidean space*, Bull. London Math. Soc. **13** (1981) 323–324.
- [53] F. Riesz and B. Sz. Nagy, **Functional Analysis**, Dover Publications, New York, 1990.
- [54] H. Nawab, T. F. Quatieri, and J. S. Lim, *Signal Reconstruction from the Short-Time Fourier Transform Magnitude*, in Proceedings of ICASSP 1984.
- [55] A. Neumaier, *Combinatorial configurations in terms of distances*, Memorandum 81-09 (Wiskunde), Eindhoven Univ. Technol., (1981).
- [56] B. Simon, **Orthogonal Polynomials on the Unit Circle. Part 2: Spectral Theory**, AMS Colloquium Publications, vol. 54, 2004.

- [57] O. Raz, N. Dudovich, O. Raz, *Vectorial Phase Retrieval of 1-D Signals*, IEEE Trans. on Signal Processing, **61**, no.7, 2013, pp. 1632–1643.
- [58] T. Strohmer and R. Heath, *Grassmannian frames with applications to coding and communications*, Appl. Comp. Harm. Anal. **14** (2003), 257–275.
- [59] J. H. van Lint and J. J. Seidel, *Equilateral point sets in elliptic geometry*, Indag. Math. **28** (1966), 335–348.
- [60] R. Vale and S. Waldron, *Tight frames and their symmetries*, Constr. Approx. **21** (2005), no. 1, 83–112.
- [61] I. Waldspurger, A. dAspremont, S. Mallat, *Phase recovery, MaxCut and complex semidefinite programming*, arXiv submission , arXiv:1206.0102
- [62] W. K. Wootters and B. D. Fields, *Optimal state-determination by mutually unbiased measurements*, Ann. Physics **191** (1989), no. 2, 363–381.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MARYLAND, COLLEGE PARK MD 20742
E-mail address, R. Balan: `rvbalan@math.umd.edu`




FIGURE 1. Mean-Square Error and CRLB bounds for $m = 800$ (top plot), $m = 600$ (middle plot), and $m = 400$ (bottom plot) when $n = 100$.







FIGURE 2. Bias and variance components of the mean-square error and CRLB bounds for $m = 800$ (top plot), $m = 600$ (middle plot), and $m = 400$ (bottom plot) when $n = 100$.

FIGURE 3. Traces for $m = 800$, $n = 100$ and $SNR = -30dB$:


 FIGURE 4. Traces for $m = 800$, $n = 100$ and $SNR = 0dB$:

FIGURE 5. Traces for $m = 800$, $n = 100$ and $SNR = 40dB$:




FIGURE 6. Mean-Square Error and CRLB bounds for $m = 800$ (top plot), $m = 600$ (middle plot), and $m = 400$ (bottom plot) when $n = 100$ and random initialization of x^0 .